

The following organizations and their boards of directors endorse the *Guidelines for Managing Substance Withdrawal in Jails: A Tool for Local Government Officials, Jail Administrators, Correctional Officers, and Health Care Professionals* published by the U.S. Department of Justice's Bureau of Justice Assistance (BJA) and National Institute of Corrections (NIC).

American Correctional Association (ACA)

American Society of Addiction Medicine (ASAM)

Foreword

Death and suffering due to withdrawal from opioids, alcohol, and other substances are preventable. Local government officials, jail administrators, correctional officers, and health care professionals have an opportunity to save lives and promote the wellbeing of individuals in jail, an opportunity bound by legal obligations set forth in the Americans with Disabilities Act and various federal civil rights acts.

To help jails and communities establish or enhance policies and procedures that appropriately address withdrawal, as well as support custody and health care staff in carrying out their responsibilities, the Bureau of Justice Assistance (BJA) and the National Institute of Corrections (NIC) are excited to present *Guidelines for Managing Substance Withdrawal in Jails: A Tool for Local Government Officials, Jail Administrators, Correctional Officers, and Health Care Professionals*. This groundbreaking document not only responds to urgent requests for guidance from the field but also advances the National Drug Control Strategy of improving access to medication for opioid use disorder for populations who are incarcerated or reentering the community.[1]

We understand that jails have a wide range of medical capabilities and encourage each facility to explore options for implementing the guidelines within their systems and communities. This will involve collaboration between jail administrators and their providers to establish policies and procedures for custody staff and health care professionals. Implementation of the guidelines also calls for collaboration within the community: namely, assessment, planning, and coordination with hospitals, local emergency medical services, opioid treatment providers, county partners, and other entities. BJA and NIC will support implementation efforts of jails and their partners through BJA's Comprehensive Opioid, Stimulant, and Substance Use Program (COSSUP) Jail Resources, offering a variety of training and technical assistance opportunities, including resources, peer-to-peer learning, and communities of practice.

We profoundly appreciate the time and commitment of the jail administrators, clinicians working in criminal justice settings, and substance use disorder specialists who wholeheartedly shared their expertise, experience, and energy with us to create these guidelines. Many of these experts are now poised to help jails and their partners implement the guidelines through training and technical assistance offered through BJA's COSSUP Jail Resources.

Thank you is also extended to the dozens of field reviewers from federal agencies and national associations who provided critical feedback on the draft document, as well as staff members from Advocates for Human Potential, Inc., the American Society of Addiction Medicine, and the National Commission on Correctional Health Care, who collaboratively led the many facets of writing and managing this effort.

Together, we can expand access to high-quality and continuous care, thereby preventing deaths from withdrawal and overdose and paving the way toward long-teessionals.

Acknowledgements

Guidelines for Managing Substance Withdrawal in Jails is the product of the vision, expertise, and perseverance of many people. The following individuals are acknowledged for their extensive contributions:

Federal Champions

- Stephen Amos, National Institute of Corrections, U.S. Department of Justice
- Margaret Chapman, Bureau of Justice Assistance, U.S. Department of Justice
- Ruby Qazilbash, Bureau of Justice Assistance, U.S. Department of Justice

Expert Committee Members

- Jeffrey Alvarez, M.D., CCHP
- Andrew F. Angelino, M.D.
- Oscar Aviles, CPM, CJM, CCE, CCHP
- Chief Deputy Jeff Clark, M.J.A.
- Kevin A. Fiscella, M.D., M.P.H., CCHP
- Carrie Hill, Esq.
- Miriam S. Komaromy, M.D.
- Shannon Robinson, M.D.
- Sandra Springer, M.D.
- Marc F. Stern, M.D., M.P.H.
- Chief Deputy Geoff Stobart
- Donna Strugar-Fritsch, B.S.N., M.P.A., CCHP

Technical Support Team Members

Advocates for Human Potential, Inc.

- Linda Frazier, M.A., RN, MCHES
- Kay S. Peavey, B.A.
- Pam Rainer, M.S.W.
- Deann Rogers, M.S.

American Society of Addiction Medicine

- Maureen Boyle, Ph.D.
- Taleen Safarian, B.A.

National Commission on Correctional Health Care

- Brent Gibson, M.D., M.P.H., CCHP-P
- Wendy Habert, M.B.A., PHR, CCHP
- Laura Lutgen-Nieves, Ph.D.
- Jim Martin, M.P.S.A., CCHP
- Deborah Ross, CCHP
- Jaime Shimkus, CCHP
- Kim Sterling, M.B.A.
- Claire Wolfe, M.A., M.P.H., CCHP

Other federal partners who contributed to this work include:

- Federal Bureau of Prisons
- National Institute on Drug Abuse
- Substance Abuse and Mental Health Services Administration
- U.S. Department of Justice, Civil Rights Division
- U.S. Department of Justice, Office of Community Oriented Policing Services
- U.S. Department of Veterans Affairs

Opioid Withdrawal 41
Screening 41
Monitoring for Withdrawal Signs and Symptoms 42
Clinical Assessment and Diagnosis 43
Medications 43
Level of Care 50
Monitoring Patients During Withdrawal Management 50
Supportive Care 51
Pregnancy and Postpartum 51
Reentry 53
Stimulant Withdrawal 55
Screening 56
Monitoring for Withdrawal Signs and Symptoms 56
Clinical Assessment and Diagnosis 56
Monitoring Patients During Withdrawal Management 57
Medications 57
Level of Care 59
Supportive Care 59
Pregnancy and Postpartum 60
Appendixes 61
Appendix A: Development Team

Task 1: Literature Review 68
Task 2: Identi cation of Draft Statements and Appropriateness/Feasibility Rating 70
Appendix C: Preferred Terminology 73
Appendix D: ASAM's Principles of Drug Testing During Withdrawal Management 75
Appendix E: Monitoring Patients During Withdrawal Management 77
Appendix F: Withdrawal Pharmacotherapy 79
Appendix G: Indicators of Dehydration 81
Appendix H: Expert Committee Members' Current Industry Relationships 83
Appendix I: Field Reviewers 85
Appendix J: Field Reviewers' Current Industry Relationships 89
Appendix K: Glossary 91
Appendix L: Acronym Glossary 97
Appendix M: Resources 99
Clinical Practice Studies 99
Legislation, Rules, and Regulations 99
Pregnancy and Postpartum 100
Substances 100
Treatment and Medication 101
Reentry 102
Screening and Assessment 103
Telehealth 104
Endnotes 105

Background

Sixty-three percent of individuals sentenced to jail (compared to 5 percent of adults in the general population) meet diagnostic criteria for SUD.[2] The typical short stays in jail, combined with scant resources for supportive services, present challenges for addressing substance withdrawal and SUD, yet jurisdictions are responsible for the health and well-being of those in custody.

Acute substance withdrawal, left unaddressed, can result in serious health complications and death. The prevalence of such deaths is difficult to determine because they often have been categorized as "illness" or "other" (e.g., aspiration pneumonia due to severe vomiting, profound dehydration) due to lack of a specific reporting category for deaths associated with drug or alcohol withdrawal.[3]

Suicide is the leading single cause of death in jails,[4] but the exact role of substance withdrawal in these deaths is difficult to quantify.[5] Risk for suicidal ideation and attempts is increased among individuals in substance withdrawal and those with an SUD. Notably, individuals with OUD have a threefold higher risk for suicidal behavior than those without OUD.[6]

Wrongful death lawsuits and other litigation for inadequate medical care are resulting in large financial settlements or judgments against counties, jail administrators, staff, and health care providers. *Managing Substance Withdrawal in Jails: A Legal Brief* provides an overview of key legislation and significant court cases related to substance withdrawal, as well as steps for jails and communities seeking to create a comprehensive response to SUD.

Beyond complying with the law, effectively managing withdrawal and SUD has significant potential for individual and societal benefits. For example, patients who received methadone treatment from a community provider prior to entering jail and continued to receive methadone from the same community provider while in jail "were less likely to receive disciplinary tickets while incarcerated and more likely to re-engage in community-based [methadone treatment] after release compared to those who underwent forced withdrawal from methadone."[7] Long-term pharmacotherapy for OUD mitigates opioid withdrawal syndrome *and* minimizes the risk of opioid overdose death, which is significant during the weeks following release from incarceration due to reduced tolerance.

Opioid Withdrawal Considerations

Effective management of opioid withdrawal involves initiation of long-term buprenorphine or methadone. When administered in a timely manner, these medications prevent withdrawal *and* treat OUD. (Naltrexone, another U.S. Food and Drug Administration-approved [FDA] medication for OUD, is not approved for withdrawal management purposes, does not relieve withdrawal symptoms, and may precipitate withdrawal.)

Opioid withdrawal management without ongoing pharmacotherapy does not treat the underlying OUD and leaves the patient at risk for overdose and death.

Development of the Guidelines

Responding to the urgent need for recognition and appropriate management of substance withdrawal, the Bureau of Justice Assistance (BJA), in partnership with the National Institute of Corrections (NIC), formed a collaboration with Advocates for Human Potential, Inc. (AHP), the American Society for Addiction Medicine (ASAM), and the National Commission on Correctional Health Care (NCCHC) to develop guidelines informed by evidence-based medical practice, while accounting for the practical realities of jails and community providers. Collectively, they facilitated the work of

These guidelines were developed using a modified RAND/UCLA Appropriateness Method (RAM). RAM is a systematic approach encompassing review of existing literature, clinical guidelines, and standards; appropriateness ratings; stakeholder comment and reconciliation; and document development.[8] The process combines scientific evidence and clinical knowledge to determine the appropriateness of a set of clinical procedures. The modified RAM used for the development of these guidelines combined the best available scientific evidence with the collective judgment of clinical and jail administration experts weighing both clinical best practices and feasibility in jail settings. (For a full description of the methodology, go to appendix B.)

Content Overview

Following a brief description of the withdrawal management process, the recommendation statements crafted by the EC and supporting narrative are presented in five sections. General Guidance addresses issues that are universal to withdrawal management in jails; the ensuing sections focus on substance-specific considerations for alcohol, sedatives, opioids, and stimulants.[a]

The numbered recommendation statements in each section provide guidance for establishing policies and procedures related to withdrawal management. Some recommendations are included to help clinicians make decisions about services and levels of care for individuals at risk for or experiencing withdrawal. Other recommendations are intended to ground jail administrators, and to some degree health directors, in the basics of withdrawal management to:

- Facilitate effective management of jail resources and foster community partnerships.
- Inform scope-of-work requirements when creating requests for contract proposals and in writing the resulting contracts in jails that outsource health care services.

It is the jurisdiction's responsibility to determine how best to apply these recommendations (e.g., whether additional staff or contracts are needed, updating policies and procedures). Recommendations in this document do not supersede any federal, state, local, or tribal regulations.

A Note on Terminology

Using clinically correct and non-stigmatizing language promotes understanding of SUD as a complex medical condition. "Withdrawal management" is used in this document to describe services to assist a patient's withdrawal from substances, a process involving far more than removing substances from the body (commonly referred to as "detoxification" or "detox"). The authors have made every effort to incorporate appropriate terminology and encourage readers to use and promote the use of person- rst and non-stigmatizing language. (For more information, go to appendix C.)

Clinical staff are purposely identified throughout this document (e.g., "physician" is used only when a physician is required), with the expectation that clinical staff meet and adhere to the requirements of their position, as well as operate within their scopes of practice. Clinical staff referred to in this document include physician, prescriber, provider, qualified health care professional, qualified health care staff, qualified mental health care professional, and responsible provider. Likewise, the terms "individual" and "patient" are used intentionally: "individual" refers to a person who is not currently being treated for substance withdrawal or SUD, whereas "patient" refers to a person whose substance withdrawal or SUD is being treated. All terms defined in the glossary are hyperlinked to their definition at first use.

[a] Nicotine withdrawal may exacerbate other withdrawal symptoms but is beyond the purview of this document. NCCHC standards include availability of nicotine replacement products when ordered by a physician.

Table G-2: Withdrawal Risk Triage by Substance

Substance	Refer for Immediate Clinical Assessment	Monitor for Withdrawal Signs or Symptoms[b]
Alcohol	• Appears unwell to a layperson. • Self-report of ≥8 standard drinks/day for men or ≥6 standard drinks/day for women, ≥4 days/week. • Reports past-week alcohol use and a history of complicated alcohol withdrawal (e.g., withdrawal-related seizures, delirium, hallucinations).	• Self-reported risk for alcohol withdrawal. • Reports recent alcohol use below the threshold specified for immediate clinical assessment AND does not report history of complicated alcohol withdrawal.

Table G-3: Assessment and Action (continued)

Consider transfer to a higher level of care when:
• History of severe or complicated withdrawal is identified. • Moderate to severe withdrawal from multiple substances is anticipated. • History of severe psychiatric symptoms is identified. • A pregnant patient may be at risk for alcohol or opioid withdrawal.
Transfer to a higher level of care when:
• Nursing, medical, or psychiatric resources recommended in this guideline are not immediately available. • Overdose is suspected. • Significant signs and symptoms persist despite multiple doses of medication. • Severe signs or symptoms develop during withdrawal management. • Existing medical or psychiatric condition worsens. • Unstable vital signs do not respond to medications. • There is severe or ongoing oversedation. • There is moderate to severe withdrawal with significant comorbidity. • Alcohol or sedative withdrawal is severe. • Complicated symptoms (seizures, delirium, hallucinations) exist. • Wernicke encephalopathy is known or suspected. • A patient cannot take required medication orally, and there is no capacity to provide medication by another route. • Severe psychiatric symptoms are present, and a mental health assessment cannot be immediately provided. • Barbiturate or gamma-hydroxybutyric acid (GHB) withdrawal is known or suspected. • The treatment plan as recommended by a qualified health care professional cannot be adequately or safely managed in the jail. • Acute medical signs or symptoms cannot be safely managed.

Recommendations

G-11. All individuals who are referred for immediate clinical assessment for substance withdrawal should be assessed by a qualified health care professional.

G-12. The initial clinical assessment should:

A. Identify any emergent medical or psychiatric needs.

B. Evaluate current withdrawal signs and symptoms.

C. Evaluate risk for severe or complicated withdrawal.

D. Assess risk for suicide.

E. Determine the appropriate level of care.

G-13. All individuals at risk for withdrawal should be assessed for SUD.

G-14. If patients are unable to engage in the full clinical assessment due to intoxication or withdrawal, they should be assessed for immediate clinical needs. The full assessment should be completed once their intoxication or withdrawal symptoms have resolved.

G-17. See substance-specific guidance regarding frequency of clinical assessments. If health care staff are not available onsite to manage the minimum required frequency of clinical assessments, the patient should be transferred to an appropriate medical facility. (Note that well-trained and supervised custody staff may monitor withdrawal signs and symptoms and administer validated tools, such as the COWS and CIWA-Ar.)

G-18. Where nursing, medical, psychiatric, or urgent care resources are recommended in these guidelines and are not immediately available onsite, the provider should be consulted to determine the necessary timing for transferring the patient to a higher level of care for further assessment and treatment.

A. If a provider is not available, the patient should be immediately transferred to a hospital.

G-19. For management of suspected substance overdose, follow the principles of first aid and cardiopulmonary resuscitation (CPR) and transfer to a local emergency department.

A. If the patient is not breathing, or their breathing is not sufficient, naloxone should be administered.

Withdrawal Management by Quali ed Health Care Professionals

Recommendations

G-20. Medical decisions should be independent of the custody level or classification of the patient.

G-21. While comprehensive clinical assessment of the patient is critical for treatment planning, treatment should not be delayed while awaiting completion of all assessments or laboratory testing or results.

G-22. Any protocols or similar guidance used in withdrawal management should reflect current clinical treatment standards and be approved by the responsible provider with input from a physician with withdrawal management expertise as appropriate.

G-23. Some patients may be released from jail prior to completion of withdrawal management. However, treatment should not be delayed based on the potential timeline for release.

G-24. A provider should receive a daily census of all patients being monitored or treated for substance withdrawal and review the status of all patients being monitored for acute withdrawal daily.

G-25. Withdrawal from alcohol, sedatives, and opioids can be dangerous and potentially life-threatening. The intensity of withdrawal cannot always be predicted. Frequent clinical assessments should be conducted by qualified health care professionals based on patient-specific orders from the provider.

A. These clinical assessments should be conducted by qualified health care professionals not less than twice per day, not more than 16 hours apart, unless otherwise stated in the substance-specific guidance in this document.

B. Unless otherwise ordered by the provider, during each clinical assessment, a qualified health care professional should evaluate:

i) General physical condition.

ii) Vital signs.

iii) Hydration.

iv) Orientation.

v) Sleep.

vi) Mental health status including risk for suicide and self-harm.

vii) Fall risk.

viii) Progression of the patient's withdrawal symptoms.

ix) Timing of the next assessment.

G-26. Withdrawal syndromes and the medications used to manage withdrawal can put patients at risk for falls. It is recommended that patients undergoing withdrawal management be assigned lower bunks and housed on lower floors.

G-27. Management of withdrawal does not constitute treatment for SUD. Jails should take proactive steps to engage patients with SUD in ongoing treatment, as described in Reentry.

Monitoring Patients During Withdrawal Management

Regularly monitoring patients helps staff detect destabilizing health, enabling providers' timely modification of treatment plans. Telehealth may be used to provide a broad range of clinical services, depending on what technologies are available and the needs of the patient. It is the provider's responsibility to determine when telehealth is or is not appropriate.

Consider the timing of monitoring to avoid interfering with the normal sleep schedule of the person being assessed, unless specifically ordered otherwise by the qualified health care professional.

Recommendations

G-28. All patients undergoing withdrawal from any substance should be monitored regularly for changes in condition. Monitoring should be conducted by qualified health care professionals or well-trained and supervised custody staff (see Staffing and Staff Training).

G-29. The onset and intensity of withdrawal is variable. The frequency and duration of monitoring should be determined based on the type of substance taken, when it was last taken, and the suspected duration of effect. Frequency of monitoring should be determined by a qualified health care professional.

A. See respective substance-specific section for monitoring intervals (see appendix E for summary of recommendations).

B. The frequency of monitoring and assessment discussed in this document is the minimum recommended for the patients described. Clinical staff should use their judgment to determine when more frequent assessment or monitoring is needed.

C. The frequency of monitoring or assessment should not be decreased or discontinued without a patient-specific order from a physician.

G-47. All custody staff should receive training that addresses the misconceptions and stigma regarding SUD, SUD treatment, and the use of medications to treat OUD.

G-48. The jail's first aid course should include training on giving CPR; managing overdose (checking respirations, positioning patient to avoid aspiration, and administering naloxone); and managing seizures (preventing head trauma, positioning the patient to avoid aspiration) while awaiting emergency medical services (EMS).

G-49. All training relevant to these guidelines should be repeated at least every 2 years.

G-50. Policy and procedures for staff training should be reviewed at least every 2 years.

Additional staff training is noted in the substance-specific sections.

Quality Assurance

Quality assurance is essential for effectively implementing these guidelines. Quality assurance should involve both custody and health care staff and include structured procedures. Ideally, meetings would take place to review data and minutes reflecting actions taken.

Recommendation

G-51. Jails should establish a quality assurance process for withdrawal management in their facility.

Supportive Care

Supportive care for withdrawal helps alleviate common physical complications and reassures the patient about what to expect from the process. For example, minimizing environmental stimulation (e.g., dimming lights, reducing noise levels) and housing patients experiencing withdrawal in smaller units (when available) may make withdrawal less physically and emotionally challenging. Isolating patients who are experiencing withdrawal, however, is not advised, due to the increased risk of self-harm (see G-63).

In addition, vomiting and diarrhea can occur during withdrawal. Dehydration and electrolyte imbalances can have serious health consequences, including death, so maintaining fluids is essential (see G-54). It is important that custody staff are alert to indicators of dehydration (see appendix G) and well-trained to report concerns about dehydration to health care staff. Qualified health care professionals will monitor for signs and symptoms of dehydration during clinical assessments.

On a broader level, supportive care encompasses all programming and resources that facilitate a patient's journey to recovery from SUD. It is important for patients to understand that treatment of withdrawal is not treatment of SUD.

The length of stay in jail will impact the nature of supportive care and may include Motivational Interviewing and other counseling, stress management strategies, education on making lifestyle changes, and mutual support/peer support programs. Of note, some jails use telehealth to provide individual and group counseling sessions (see Telehealth Resources).[9]

Recommendations

G-52. All patients experiencing withdrawal, regardless of severity, should receive supportive care, which may include

G-53. Patient education regarding the withdrawal process is a necessary component of treatment. Patients experiencing withdrawal should be counseled on:

A. The signs and symptoms of withdrawal.

B. The assessment process.

C. What to expect during treatment and monitoring.

D. What to do if they are released prior to completion of withdrawal management.

E. The patient's role in helping to manage withdrawal (e.g., staying hydrated, communicating with health care and custody staff).

F. The importance of engaging in ongoing SUD treatment to support sustained recovery.

G-54. Unless restricted access is ordered by a provider and the patient is placed under close medical supervision, patients experiencing withdrawal should have unimpeded access to water or electrolyte solution.

A. Both health care and custody staff should encourage patients to stay well-hydrated.

B. Custody staff should notify health care staff if they notice that a patient undergoing withdrawal management has insufficient fluid intake or is refusing to drink.

Reentry

Reentering the community is a challenging time for individuals, especially for those with SUD. Research shows that risk of overdose in the 2 weeks following release from incarceration is extremely high.[10] Patient education on the risks associated with reduced tolerance to substances as a result of withdrawal and how this might lead to overdose and death is important for mitigating these risks. This topic is further discussed in Opioid Withdrawal.

Ideally, individualized reentry plans that address ongoing withdrawal management needs, SUD treatment engagement, overdose prevention, and recovery supports are developed with appropriate input from the health care provider for all patients with SUD. These plans include assertive referrals (i.e., scheduling appointments for community care), as well as strategies for:

- Appropriate sharing of health records (electronic or paper) as authorized by state law.
- Establishing insurance coverage including support for enrollment or re-activation of suspended or terminated Medicaid coverage (if applicable).
- Arranging patient navigator support or collaborating with community substance use treatment, such as providers

C. Jails should have policies and procedures that support continuity of care through sharing of health records and clinical information with community providers.

Suicide

appropriate, obtain informed consent prior to conducting any health care-related drug testing. The intent of the notification is to ensure that the patient understands the requirements without conveying judgment.

Withdrawal during pregnancy is also associated with high rates of return to use,[12] which can lead to harm to the fetus, including the return to alcohol use that can result in fetal alcohol spectrum disorders (FASD), overdoses, and other harms. Efforts should be made to engage all pregnant patients treated for withdrawal in ongoing SUD treatment. For pregnant patients with OUD, the standard of care is ongoing treatment with buprenorphine or methadone.

Jails should establish policies and protocols related to withdrawal management during pregnancy. These policies and protocols should be developed and approved by a physician or physicians with experience in SUD and obstetrics and communicated to all facility and community decision-makers, from arrest through reentry.

Additionally, jails are encouraged to establish policies and procedures for distributing naloxone or providing prescriptions for naloxone to patients, including those who are pregnant or postpartum, as they leave custody to reenter the community.

Recommendations

G-64. All patients of childbearing potential or childbearing age should be assessed for pregnancy and offered pregnancy testing.

G-65. Health care professionals and custody staff should not assume symptoms such as nausea, headache, anxiety, and insomnia are due to pregnancy and should remain vigilant for substance withdrawal.

G-66. When offering medications to pregnant or lactating patients to treat substance withdrawal or medical complications of withdrawal, the prescriber should discuss the risks and benefits of each medication for the patient and the fetus or infant, as well as risks associated with untreated withdrawal and ongoing substance use during pregnancy.

G-67. Initiation and continuation of medication should not be withheld or delayed due to a patient's pregnancy or lactation status.

G-68. Health care professionals should follow the recommendations included in the substance-specific sections on treating withdrawal in pregnant patients.

G-69. Some states have laws that require health care providers to report positive drug test results and impose adverse legal and social consequences on individuals who use substances during pregnancy. Therefore, qualified health care professionals should educate pregnant patients regarding these risks and obtain informed consent prior to conducting any health care-related drug testing.

Older Adults

Approximately 8 percent of adults confined in local jails are age 55 and older,[13] a cohort referred to in this document as "older adults." Fifty-five years is used in this context because heavy substance use, inadequate health care, and the stress of incarceration itself contribute to accelerated aging, which means that individuals may have physical and mental conditions that are typically associated with those who are at least 10 years older.[14] Hypertension, diabetes, angina, heart attacks, arthritis, and hepatitis are more prevalent among individuals in jail custody who are ages 50–65 than those in the same age group who are not incarcerated.[15]

Withdrawal management in older adults can be complicated by the higher prevalence of chronic diseases that impact

A-7. Alcohol withdrawal severity assessment scales (including the CIWA-Ar) should not be used as stand-alone diagnostic tools because scores can be influenced by conditions other than alcohol withdrawal.

A-8. Alcohol withdrawal can progress rapidly with serious health consequences. Individuals referred for immediate clinical assessment should be assessed immediately by a qualified health care professional to determine their risk for developing severe and/or complicated alcohol withdrawal or complications from alcohol withdrawal.

A. In addition to signs and symptoms of alcohol withdrawal, the following factors are associated with increased patient risk for complicated withdrawal or complications of withdrawal and should be assessed:

i. History of complicated alcohol withdrawal (e.g., delirium or alcohol withdrawal seizure).

ii. Multiple prior withdrawal episodes.

iii. Comorbid illnesses (especially traumatic brain injury).

iv. Age > 65.

v. Marked autonomic instability on presentation.

vi. Long duration of heavy and regular alcohol consumption.

vii. Seizure(s) during the current withdrawal episode.

viii. Physiological dependence on sedatives such as benzodiazepines or barbiturates.

ix. Positive blood alcohol concentration in the presence of signs or symptoms of withdrawal.

A-9. If a toxicology test (blood, breath, or urine) for alcohol use is used, do not rule out the risk of developing alcohol withdrawal if the test result is negative.

Withdrawal Management by Quali ed Health Care Professionals

Historically, alcohol has been administered to prevent withdrawal in some cases. However, administration of oral or IV alcohol has no proven efficacy, no accepted protocols, and can be toxic.

Recommendations

A-10. Oral or IV alcohol should not be used for the prevention or treatment of alcohol withdrawal.

Monitoring Patients During Withdrawal Management

During withdrawal management, monitoring typically includes both collection of vital signs and administration of a validated severity assessment scale, such as the CIWA-Ar (see table A-2).

Table A-2: Alcohol Withdrawal Severity [19]

Severity Category	Mild	Moderate	Severe	Complicated
Associated CIWA-Ar Range	< 10	10–18	≥ 19	≥ 19

As alcohol withdrawal is primarily treated with benzodiazepines, which are sedative medications, it is important that staff monitor for and report oversedation.

Recommendations

A-11. Alcohol withdrawal severity should be monitored using a validated tool, such as the CIWA-Ar.

A-12. Use of the CIWA-Ar does not replace a clinical assessment, including collection of the patient's vital signs.

A-13. A clinical assessment including the CIWA-Ar should be conducted at least every 8 hours during alcohol withdrawal treatment until the CIWA-Ar score remains below 10 for 24 hours.

A. If the CIWA-Ar is > 19, repeat the CIWA-Ar at least every 6 hours during alcohol withdrawal treatment until the score falls below 19, then continue monitoring with the CIWA-Ar at least every 8 hours until the score remains below 10 for 24 hours.

B. More frequent monitoring should be conducted if clinically indicated.

A-14. To support management of alcohol withdrawal, a qualified health care professional, or custody staff, who has been well-trained to administer the CIWA-Ar (or another validated tool) should be available at all times.

A-15. Custody staff should be well-trained to identify oversedation and to alert health care staff if a patient appears oversedated.

A-16. Regular clinical assessments should monitor for dehydration. Health care staff should be alerted if the patient reports, or custody staff otherwise become aware, that the patient has stopped drinking, has a significant reduction in urine volume or frequency of urination, or has very dark urine.

Level of Care

Severe alcohol withdrawal can result in serious health consequences, including death. The clinical assessment should inform the determination of whether the predicted severity of withdrawal can be managed in the jail or transfer to a hospital is indicated.

Table A-3: Signs and Symptoms of Wernicke Encephalopathy–Korsakoff Syndrome [20]

Wernicke Encephalopathy	Korsakoff Syndrome
• Confusion or loss of mental activity. • Loss of muscle coordination (ataxia), manifested as leg tremor or wide-based or unsteady gait. • Vision changes: o Abnormal eye movements (back and forth movements called nystagmus). o Double vision.	• Inability to form new memories. • Loss of memory, can be severe. • Making up stories (confabulation).

Recommendations

A-17. Level-of-care determination should be based on a patient's risk for developing severe or complicated alcohol withdrawal, or complications of withdrawal, as well as current signs and symptoms.

A-18. If the patient requires transfer to a higher level of care, treatment for immediate needs should be initiated while awaiting transfer, where feasible.

A-19. While most patients experiencing alcohol withdrawal may be managed in the facility, if the following indications are present, the patient should be transferred to a hospital (unless the jail has hospital-level capacity):

A. Agitation or severe tremor persists despite having received multiple doses of medication.

B. Severe signs or symptoms, such as persistent vomiting, marked agitation, hallucinations, confusion, or seizure, develop.

C. Existing medical or psychiatric condition worsens.

D.

Dosage Regimens

Front loading: An approach where moderate to high doses of a long-acting medication are given frequently at the start of treatment to achieve rapid control of withdrawal signs and symptoms. Front loading can be followed by a symptom-triggered or fixed-dose regimen.

Fixed dosing: An approach where a predetermined dose (which can be determined based on withdrawal severity) is administered at fixed intervals according to a schedule. Doses usually decrease in a gradual taper over several days. Additional medication may be provided if the fixed dose does not adequately control symptoms.

Symptom-triggered dosing: An approach where patients are given medication only when symptoms cross a threshold of severity (e.g., 15 mg oxazepam for CIWA-Ar scores 8–15, 30 mg oxazepam for CIWA-Ar > 15).

WARNING: Symptom-triggered dosing, if used alone, requires highly reliable and dedicated staff to avoid undertreatment of withdrawal and should be used only when there are adequate qualified staff to perform assessments.

Recommendations

A-24. Benzodiazepines are the preferred agent for treating alcohol withdrawal.

A-25. While no particular benzodiazepine agent is more effective than another, long-acting benzodiazepines are the preferred agents due to the clinical benefits of their longer duration of action. Adequate treatment with a long-acting benzodiazepine is effective in preventing withdrawal seizures.

A. See *The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management* for more detailed clinical guidance and dosing protocols, including guidance regarding patients with severe liver disease and other significant comorbidities.

A-26. Patients with CIWA-Ar scores < 10 and who are at minimal risk of developing severe or complicated alcohol withdrawal may be provided supportive care alone and monitored.

A. It is also appropriate to use benzodiazepines prophylactically for alcohol withdrawal.

A-27. Patients with CIWA-Ar scores > 10 and patients at risk of developing severe or complicated alcohol withdrawal should receive pharmacotherapy and supportive care.

A-28. Front loading, fixed dosing, and symptom-triggered dosing are all appropriate in jail settings and may be used in combination.

A-29. Front loading is recommended for patients at risk for severe or complicated alcohol withdrawal (e.g., CIWA-Ar ≥ 21). Diazepam and chlordiazepoxide are preferred agents for front loading.

A-30. The patient's signs and symptoms should be monitored, as discussed in Monitoring Patients During Withdrawal Management in General Guidance, regardless of the dosing schedule used.

A-31. Qualified health care staff should monitor patients taking benzodiazepines for signs of oversedation and respiratory depression.

A-32. If a patient's symptoms are not controlled as expected with the dose of benzodiazepine prescribed, the provider should consider increasing the dose.

A-33. Benzodiazepines used to treat alcohol withdrawal should be tapered and discontinued following treatment.

A-34. If the patient has been using sedative substances and alcohol, withdrawal should be handled in the same way as withdrawal from sedatives (see recommendations in Sedative Withdrawal).

A-35. Do not give antiseizure medications prophylactically to prevent alcohol withdrawal unless the patient also has an underlying seizure disorder.

A-36. Alcohol use disorder (AUD) is a chronic medical condition. Patients should be provided with information about the range of evidence-based treatments, including medications for AUD, and recovery support services available in the community.

A. The provider should offer to initiate evidence-based treatment for AUD, including medications and regular follow-up visits at clinically appropriate intervals. Follow-up visits may include monitoring for return to use, Motivational Interviewing to encourage engagement in AUD treatment, psychoeducation, and management of medications for AUD.

A-37. If a patient entering jail is taking prescribed medication for AUD, the medication should be continued unless otherwise ordered by a prescriber based on documented clinical need.

Supportive Care

Inadequate nutrition associated with heavy alcohol use can lead to conditions that can typically be prevented through supportive care. For example, thiamine can reduce the risk for Wernicke encephalopathy.

Recommendations

A-38. As discussed in General Guidance, supportive care is appropriate for all levels of alcohol withdrawal severity and may include nutritional supplementation, IV fluids, glucose, management of electrolyte abnormalities, and periodic clinical reassessments, as clinically indicated (see Monitoring Patients During Withdrawal Management).

A. If dehydration is an ongoing concern and the jail does not have the capacity to manage IV fluids, the patient should be transferred to a facility that does.

A-39. Alcohol use can be associated with vitamin and mineral deficiencies.

A. To reduce the risk of encephalopathy, all patients with suspected AUD should be treated with thiamine, 100 mg daily, either orally or intramuscularly for 3–5 days.

B. Other vitamin and mineral supplementation (e.g., magnesium, folate) should be considered, as determined by the treating provider.

Pregnancy and Postpartum

Alcohol withdrawal presents significant risks during pregnancy, often requiring care at a hospital with expertise in managing high risk obstetrical cases and with fetal monitoring available.

Recommendations

A-40. Pregnant patients should typically be transferred to a setting with 24-hour medical care available (e.g., emergency department, hospital).

A-41. If managing the patient's alcohol withdrawal in the jail, consult with a provider experienced in obstetrical care.

A. Specialty consultation should not delay initiation or titration of treatment.

A-42. Benzodiazepines are the preferred medication for treatment of pregnant patients with alcohol withdrawal. While there is a risk of teratogenicity during the first trimester, the risk appears small and balanced in view of the risk for FASD and consequences to the mother and fetus should severe maternal alcohol withdrawal develop.

A-43. For patients at risk for preterm delivery or in the late third trimester, use of a short-acting benzodiazepine is recommended. This minimizes the risk for neonatal benzodiazepine sedation given shorter onset and duration of action.

A-44. Given the risk of FASD, pregnant patients with alcohol withdrawal should be educated on the importance of treatment for AUD.

A-45. Pregnant patients should be informed of all wraparound services that will assist them in addressing newborn needs (including food, shelter, and pediatric clinics for inoculations) and programs that will help with developmental or physical issues that the newborn may experience as a result of in-utero alcohol exposure.

A-46. Qualified health care professionals should understand and follow their state laws regarding definitions of child abuse and neglect, reporting requirements, and plans of safe care for newborns with in-utero alcohol exposure.

Managing Comorbidities

Co-occurring medical and psychiatric conditions can complicate the management of alcohol withdrawal and should be considered when developing the treatment plan.

Recommendations

A-47. For patients with medical comorbidities, consult with specialists as needed to modify the medication and/or protocol used for treating alcohol withdrawal.

A-48. Hallucinations that develop in the context of alcohol withdrawal may indicate an alcohol-induced psychotic disorder. A patient who develops hallucinations should be transferred to a higher level of care for treatment.

A. Benzodiazepine medication should be given concurrently with the call for transport, if feasible.

B. Antipsychotic medication can be considered, concurrent with the call for transport, if feasible and needed to manage severe psychotic sympt1bMC 1820 >>BDC(LUS)/MCID 18l for transport, if feasible.e > BD <</Lang (der)93te

Sedative Withdrawal

Table SH-1: Development of Sedative Withdrawal Symptoms

Duration of Action	Examples	Timeframe
Short-acting	alprazolam, lorazepam	First 24 hours after cessation of use
Long-acting	clonazepam, diazepam	24–48 hours after cessation of use

Some indicators of sedative withdrawal (table SH-2) may be misattributed to other causes, highlighting the importance of referring any individual who appears unwell for immediate clinical assessment.

Table SH-2: Possible Indicators of Sedative Withdrawal* [24,25,26]

- Abdominal cramping, nausea, and vomiting
- Anxiety
- Circulatory failure
- Delirium (including disorientation and hallucinations)
- Hyperthermia and sweating
- Increased heart rate and blood pressure
- Insomnia
- Occasional psychotic thoughts
- Restlessness
- Seizures
- Tremors
- Unease

*Staff should be alert to any indicators that the individual is unwell, not only those listed here.

Recommendations

SH-3. Individuals who self-report risk for sedative withdrawal and those who report sedative use below the threshold specified in SH-1 (daily or near daily and within past 7 days) should be monitored at least every 6 hours for at least the first week for emergence of withdrawal signs and symptoms.

SH-4. Although a few benzodiazepine withdrawal scales have been developed, there is no single scale that has been well validated and replicated. As discussed in General Guidance, if signs or symptoms emerge or if the individual begins to appear unwell to a layperson, they should be referred for immediate clinical assessment.

Clinical Assessment and Diagnosis

All sedative withdrawal should be treated, regardless of length of jail stay. Clinical assessment will determine the type and intensity of care required to address the patient's needs.

No validated scales for assessing sedative withdrawal exist. The Clinical Institute Withdrawal Assessment Scale – Benzodiazepines (CIWA-B) is a 22-item instrument for assessing and monitoring symptoms of benzodiazepine withdrawal, but its validity and reliability have not been fully determined.[27]

Recommendations

SH-5. As with all withdrawal, a patient history and physical exam including vital signs should be done to assess withdrawal severity and inform treatment.

SH-6. Do not use the CIWA-Ar for assessing sedative withdrawal.

SH-7. Because of the high risk of delirium, seizures, and death, sedative withdrawal should always be treated.

Level of Care

Withdrawal from barbiturates and GHB is complex and requires hospital-level care, as does severe or complicated withdrawal from benzodiazepines. Patients should be transferred to a hospital, preferably (when possible) a hospital with expertise in treating sedative withdrawal. Use of GHB is typically determined by patient history due to its short-acting nature. If the patient is intoxicated, obtaining this history may be difficult; however, witnesses (law enforcement, family, friends) may be able to provide information about GHB use.

Recommendations

SH-8. For sedative withdrawal, hospitalization is suggested for patients showing signs of severe or complicated withdrawal.

A. These signs may include:

i) Delirium.

ii) Hallucinations.

iii) Changes in consciousness.

iv) Profound agitation.

v) Autonomic instability (when heart rate, blood pressure, and sweating and other nonvoluntary body functions fluctuate).

vi) Seizures.

B. For patients experiencing less severe sedative withdrawal who have a history of complicated sedative withdrawal symptoms, transfer to a setting with 24-hour medical care should be considered.

C. For patients at risk for sedative withdrawal (but not presenting with signs or symptoms) who have a history of complicated withdrawal symptoms, more frequent monitoring should be considered.

SH-9. Hospitalization is recommended for patients experiencing barbiturate withdrawal, unless the jail has hospital-level capacity (i.e., telemetry [including cardiac monitoring] and full-code response including intubation/ventilation support).

SH-10. GHB withdrawal is complex and technically more difficult than other sedative withdrawal. Transfer to a hospital is recommended. When possible, the patient should be transferred to a hospital with experience treating complex sedative withdrawal.

Monitoring Patients During Withdrawal Management

Recommendations

SH-11. Patients treated for sedative withdrawal should have a daily clinical assessment by a qualified health care professional for at least the first week, or as their condition indicates.

A. After the first week, patients should be assessed by a qualified health care professional at least two times per week until the taper is complete (see SH-14).

SH-19. Do not give antiseizure medications prophylactically to prevent sedative withdrawal unless the patient also has an underlying seizure disorder.

Supportive Care

Sedative withdrawal can cause a number of side effects, including insomnia and anxiety, that may be relieved through supportive care (when possible).

Recommendations

SH-20. In addition to the recommendations in General Guidance, supportive care for patients in sedative withdrawal may include cognitive behavioral therapy (except during acute withdrawal), stress management, sleep hygiene, and relaxation training, which may be helpful both during and after taper.

Pregnancy and Postpartum

Treating sedative withdrawal can be safely managed in pregnancy, following the guidelines described above.

Recommendations

SH-21. Pregnant patients should undergo withdrawal management slowly and in consultation with an obstetrician.

SH-22. The principles of withdrawal management from sedatives are the same for pregnant and non-pregnant patients.

Opioid Withdrawal

Opioids include illicit drugs (e.g., heroin, illicitly manufactured fentanyl, and its analogues) and prescription pain relievers, such as oxycodone (Percocet or Oxycontin), hydrocodone (Vicodin), methadone, and morphine. Individuals who are physiologically dependent on opioids, including those who regularly use either illicit or prescription opioids, are at risk for opioid withdrawal.

Dangerous Changes in Drug Composition

Individuals may not always know that they have been using opioids or which opioids they have been using. Increasingly, individuals are consuming counterfeit pills or drugs containing fentanyl.[28,29] Given the high potency of fentanyl mixed in with the drug supply, multiple doses of naloxone may be necessary to reverse an overdose.[30]

It is important for qualified health care professionals to understand the regional drug supply, remain current on clinical guidance, and regularly revisit response protocols. For example, the veterinary tranquilizer xylazine, which is currently added to fentanyl and other illicit opioids, has been linked to many overdose deaths in the Northeast. Naloxone may be less effective in cases of overdoses involving opioids mixed with xylazine.[31]

Opio id withdrawal syndrome can be medically complex and, in the absence of appropriate management, life-threatening.

Recommendation

O-1. Anyone who reports regular use of opioids (including prescription opioid misuse), OUD, or opioid withdrawal risk should be considered at risk for opioid withdrawal. [Note that this does not apply to use of prescription opioids taken as prescribed, as the patient should have continued access to the medication.]

A. If prescription opioids are discontinued, based on a patient-specific order from a provider (see G-4), the patient should be considered at risk for opioid withdrawal.

Monitoring for Withdrawal Signs and Symptoms

Opioid withdrawal syndrome refers to the wide range of symptoms that occur when individuals who regularly take opioids, either illicit or prescribed, stop or reduce their use. Regular monitoring for signs and symptoms of withdrawal supports rapid identification and referral for clinical assessment.

Some individuals may initially screen negative for withdrawal risk but later exhibit signs or symptoms of withdrawal (tables O-1 and O-2). Custody staff should be well-trained to refer all individuals who appear unwell for immediate clinical assessment (see G-44), regardless of initial screening results.

Table O-1: Opioid Withdrawal Symptom Emergence and Duration [33]

	Emergence	**Duration**
Short-acting opioids (e.g., heroin, oxycodone)	Within 12 hours of last use and peaking within 24–48 hours	3–5 days
Long-acting opioids (e.g., methadone)	Within 30 hours of last use	Up to 10 days

Table O-2: Possible Indicators of Opioid Withdrawal* [34]

- Agitation
- Anxiety
- Abdominal cramping
- Diarrhea
- Dilated pupils
- Increased tearing
- Insomnia
- Muscle aches
- Nausea
- Piloerection (“goose bumps”)
- Runny nose
- Sweating
- Vomiting
- Yawning

* Staff should be alert to any indicators that the individual is unwell, not only those listed here.

Recommendations

O-2. Individuals may be considered at risk for opioid withdrawal even if they do not meet the clinical criteria for an OUD.

O-3. Individuals who report regular opioid use (including prescription opioid misuse) or screen positive for current OUD should be monitored for signs and symptoms of opioid withdrawal at least every 4 hours for the first 72 hours of incarceration.

A. As discussed in General Guidance, individuals at risk for withdrawal should ideally be housed together in a dedicated unit to facilitate monitoring.

B. Patients who appear unwell to a layperson or score ≥ 3 on the COWS should be referred for immediate clinical assessment.

Clinical Assessment and Diagnosis

The initial clinical assessment, conducted by a qualified health care professional, focuses on identifying any medical risks that would necessitate a higher level of care or a full medical evaluation. Validated assessment tools, such as the COWS, allow for methodical collection of information about withdrawal signs and symptoms at assessment and during monitoring.

Recommendations

O-4. The initial clinical assessment should focus on identifying signs and symptoms of opioid withdrawal, as well as overdose and withdrawal risk.

A. Assessment of withdrawal risk should focus on the types of opioids used, route of use, length of time used,

WARNING

- Concurrent use of methadone or buprenorphine with alcohol or benzodiazepines increases respiratory depression risk. However, concurrent treatment with benzodiazepines (e.g., for co-occurring alcohol or sedative withdrawal) should not be a reason to withhold treatment with buprenorphine or methadone. Increased monitoring may be appropriate.
- Naltrexone is not approved by the FDA for withdrawal management purposes and does not relieve withdrawal symptoms. It cannot be administered until patients are free of short-acting opioids for at least 7–10 days or 10–14 days for long-acting opioids. Failing to observe these wait periods may cause precipitated withdrawal.
- Anesthesia-assisted opioid detoxification or ultra-rapid opioid detoxification, which involves large doses of naloxone, anesthesia, and a diuretic, has been associated with serious complications (including cardiac arrest and death) and is not recommended.

Reco

O-15. If individuals entering jail are experiencing symptoms of moderate to severe opioid withdrawal, they should be referred for immediate clinical assessment by a provider who can initiate buprenorphine or methadone or transferred to a facility that can provide buprenorphine or methadone if indicated.

O-16. Even if the patient may not be able to maintain treatment in the community, buprenorphine or methadone should still be initiated.

O-17. Patients undergoing withdrawal management should be advised of the risk of returning to use, overdose, and overdose death if they choose not to engage in ongoing medication treatment.

O-18. Jails that do not have prescribers onsite 24 hours per day, 7 days per week should consider using telehealth to support access to buprenorphine and methadone treatment.

O-19. Opioid withdrawal management is not necessary if the patient is immediately initiated on methadone or buprenorphine upon intake to the jail. However, if opioid withdrawal management is clinically indicated or if the patient declines ongoing treatment with medications for OUD, buprenorphine or methadone are also recommended for withdrawal management (see *The ASAM NPG for the Treatment of OUD* for tapering guidance).

O-20. For protocols for initiating medications for OUD, refer to *The ASAM NPG for the Treatment of OUD*.

O-21. Jails should establish mechanisms for rapidly initiating methadone or buprenorphine treatment when the provider is off-site.

O-22. Adjunctive medications (e.g., clonidine, anti-nausea medication) should not be necessary if buprenorphine or methadone is provided in adequate doses (typically ≥ 16 mg/day of buprenorphine; see *The ASAM NPG for the Treatment of OUD* for methadone-dosing recommendations) and should not be used in the place of an adequate dosage of buprenorphine or methadone.

A. Patient discomfort during opioid withdrawal management may indicate the dose of buprenorphine or methadone is too low.

O-23.

Buprenorphine

Buprenorphine is an effective medication for treating both opioid withdrawal and OUD. Recent changes in federal legislation allow buprenorphine to be prescribed by any prescriber with a DEA registration to prescribe controlled substances. However, some states may limit buprenorphine-prescribing authority.

Buprenorphine should be initiated when the patient shows objective signs or symptoms of withdrawal (e.g., pupil dilation, goose bumps, gastrointestinal discomfort). In a patient at risk for opioid withdrawal, if buprenorphine is

Table O-3: Buprenorphine Formulations (continued)

Name	Route of Administration (Dosing)	For the Treatment of	Comments
Generic: Buprenorphine extended-release Brand: Sublocade	Extended-release injection (monthly)	Moderate to severe OUD in patients who have initiated treatment with transmucosal forms, followed by dose adjustment for a minimum of 7 days.	• No risk for patient diversion or misuse. • Requires patients to be on a stable dose of transmucosal buprenorphine for at least 7 days. • Less fluctuation in buprenorphine levels (compared to daily doses).
Generic: Buprenorphine extended-release Brand: Brixadi	Extended-release injection (weekly or monthly)	Moderate to severe OUD in patients after a single dose of transmucosal buprenorphine or who are already being treated with buprenorphine.	• No risk for patient diversion or misuse. • Only a single prior dose of transmucosal buprenorphine required prior to initiation. • Less fluctuation in buprenorphine levels (compared to daily doses).

Recommendation

O-26. Buprenorphine should be initiated when the patient shows objective signs of opioid withdrawal.

A. The patient should be monitored for precipitated withdrawal for at least 30 minutes after the initial administration of buprenorphine.[c]

B. If a patient is experiencing precipitated withdrawal and a prescriber is not immediately available to conduct an assessment, the patient should be transferred to a hospital.

Methadone

Methadone, a long-acting full opioid agonist, is an effective treatment for opioid withdrawal syndrome and OUD. With the exception of access under the 72-hour emergency rule, methadone for the treatment of OUD is currently only available through federally registered opioid treatment programs (OTPs).

72-Hour Emergency Rule

Under this rule, a physician outside of an OTP is allowed to administer (but not prescribe) methadone to a patient to relieve acute withdrawal symptoms while arranging for the patient's referral for treatment, for up to 72 hours. This may not be renewed or extended, nor may more than 1 day's medication may be administered or given to a patient at one time.[39]

Jails can either become a certified OTP[40] or establish an MOU with an external methadone treatment provider detailing roles, responsibilities, payment, and security concerns, such as:

- Transport of patients to the methadone treatment provider.
- Transport of methadone to the jail by methadone treatment provider, jail medical, or jail custody staff.
- Guest dosing for patients who are established with a different methadone treatment provider.

As noted above, properly registered mobile narcotic treatment providers provide another treatment option for correctional facilities, which may be especially useful in rural communities.

---

[c] Note: Buprenorphine may be initiated prior to the emergence of objective signs of withdrawal when using a low-dose buprenorphine initiation protocol. See ASAM's "Treatment of Opioid Use Disorder with Buprenorphine: Clinical Consideration for Treatment of Individuals with OUD using High Potency Opioids" (submitted March 23).

Methadone can be difficult to access, and discontinuation of this medication puts the patient at risk of returning to use, so reentry planning upon initiation of methadone is essential.

Table O-4: Methadone Formulations

Recommendations

O-31. It is not appropriate to require a change in or discontinuation of OUD medication for nonclinical reasons (e.g., conditions of custody, release).

O-32. Discontinuing OUD medication puts the individual at risk for opioid withdrawal, and increases the risk for returning to use, overdose, and overdose death. Jails should continue treatment for patients who, upon arrival, are taking medication for treatment for OUD. (See also recommendation statement G-4 and supporting narrative.)

O-33. Patients should not be required to transition from opioid agonist (methadone or buprenorphine) to opioid antagonist (naltrexone) treatment.

O-34. In the absence of a methadone treatment provider or a provider authorized to prescribe buprenorphine (in states that restrict buprenorphine prescribing), jails should use the 72-hour emergency prescribing authority to provide methadone or state emergency access provisions to provide buprenorphine while they make arrangements for ongoing treatment.

A. The 72-hour emergency prescribing authority (and, typically, other emergency access provisions) cannot be renewed during the individual's stay in jail and should not be used in place of establishing access to buprenorphine and methadone.

O-35. Transitioning patients currently in methadone treatment to buprenorphine is clinically complex and should be managed by, or in consultation with, a provider experienced in managing this transition.

O-36. Discontinuation of methadone or buprenorphine is clinically complex and should be done only when clinically indicated and by a medical provider with SUD treatment expertise.

O-37. Tapering methadone takes longer in patients who are in methadone treatment compared to those initiated on methadone to treat withdrawal. Jails should not attempt to manage this process without the oversight of a methadone treatment provider.

Alpha-2 Adrenergic Agonists (Second-line Treatment)

Alpha-2 adrenergic agonists (FDA-approved lofexidine and off-label use of clonidine) can be used to treat withdrawal when patients choose to taper off buprenorphine or methadone, and in preparation for initiation of extended-release naltrexone. These medications are a second-line option, because they do not relieve opioid withdrawal symptoms as effectively as buprenorphine and methadone and can cause low blood pressure. This risk is increased when patients are not adequately hydrated. In addition, there can be dangerous elevations in blood pressure when this medication is stopped in patients with existing hypertension. When used, these medications are often combined with other medications targeting specific opioid withdrawal symptoms, such as benzodiazepines for anxiety, loperamide or bismuth salicylate for diarrhea, acetaminophen or nonsteroidal anti-inflammatory medications (NSAIDs) for pain, various medications for insomnia, and ondansetron for nausea.

Recommendations

O-38. Lofexidine or clonidine may also be appropriate to manage opioid withdrawal in the rare situations where complete opioid withdrawal management is clinically indicated, including when the patient declines treatment with buprenorphine or methadone.

A. Alpha-2 adrenergic agonists are contraindicated in patients with low blood pressure. As these medications can cause hypotension, blood pressure should be monitored regularly in patients prescribed these medications by personnel well-trained in blood pressure measurement.

B. See *The ASAM NPG for the Treatment of OUD* and SAMHSA's *Medication-Assisted Treatment for Opioid Addiction in Opioid Treatment Programs, Treatment Improvement Protocol (TIP) 43)* for dosage recommendations.

Level of Care

As discussed in General Guidance, the level of care should be appropriate for the anticipated severity of the withdrawal syndrome, as well as any medical or psychiatric comorbidities present. Where treatment services and resources are recommended in this section, including buprenorphine and methadone treatment, and are not immediately available in the jail, timely transfer to a higher level of care is indicated.

Monitoring Patients During Withdrawal Management

The withdrawal severity score should gradually decrease following medication initiation. An increasing score following buprenorphine initiation may indicate precipitated withdrawal and should trigger an immediate medical assessment. As mentioned in G-33, monitoring should only be discontinued upon a patient-specific order from the prescriber based on clinical evaluation and determination that it is safe to do so.

Recommendations

O-39. Opioid withdrawal symptom severity should be monitored with a validated tool, such as the COWS.

O-40. To support management of opioid use, it is recommended that jails, at minimum, have:

A. Staff who are well-trained to administer the COWS (or another validated tool) available at all times.

B. A provider authorized to prescribe buprenorphine on staff, under contract, or under a memorandum of understanding.

C. When feasible, direct affiliation with a methadone treatment provider to provide access to methadone.

O-41. Custody staff may be trained to effectively administer the COWS, including collection of vital signs.

O-42. The onset and severity of opioid withdrawal is dependent on the type of drug taken, when it was last taken, and how long it lasts in the person's body. Monitoring intervals should be determined by a qualified health care professional based on the anticipated timing and severity of withdrawal for the individual patient.

A. Patients who report use of a short-acting opioid (e.g., heroin, oxycodone, fentanyl) should be monitored using the COWS at least every 4 hours.

B. Patients who report using long-acting opioids (e.g., extended-release formulations, methadone) should be monitored using the COWS at least every 8 hours.

O-43. In patients with concurrent sedative use disorder or withdrawal, methadone or buprenorphine should be used to stabilize withdrawal from opioids.

Recommendations

O-48. Psychosocial treatment alone or withdrawal management alone are not appropriate for pregnant or postpartum patients with OUD. Pregnant and postpartum patients who are physically dependent on opioids should receive ongoing treatment with methadone or buprenorphine to prevent harm to the patient and fetus.

O-49. Methadone or buprenorphine treatment should be initiated as early as possible during pregnancy since opioid withdrawal can increase the risk of miscarriage or premature delivery.

A. Transfer to the hospital is likely to be needed for appropriate management, especially in the third trimester of pregnancy.

O-50. Opioid withdrawal risk in pregnant patients should be treated with urgency. If the jail does not have the capacity

Reentry

Most patients who require opioid withdrawal management will have OUD. Numerous studies have demonstrated that treatment with medications for OUD during incarceration and reentry significantly reduces the risk for overdose and overdose death.[49] These studies also found that treatment for OUD with medication reduces illicit opioid use, increases engagement in OUD treatment in the community, and reduces risks of reincarceration.

If methadone or buprenorphine are abruptly reduced or discontinued upon reentry, patients will be at risk of withdrawal, as well as at risk for overdose and death.

Recommendations

O-59. Individuals who are being treated with buprenorphine or methadone should continue the medication while in jail, and the jail should assist with transfer to community-based treatment upon release.

O-60. Qualified health care professionals should ensure the patient has access to an adequate supply of buprenorphine or methadone to prevent interruption of dosing when the patient transitions to the community.

A. Backup plans are vital in the event the community appointment cannot be completed. Bridging clinics and telehealth can be very helpful.

O-61. Naloxone or a prescription for naloxone should be made available to all patients with OUD upon release.

A. Jails should consider providing naloxone or a prescription for naloxone to all patients with SUD.

B. Jails should consider making naloxone or a prescription for naloxone available to friends and family members of patients with SUD.

Stimulant Withdrawal

Approximately 10 percent of the criminal justice population surveyed used methamphetamine or cocaine or misused prescription stimulants (e.g., nonmedical use of amphetamine products for treating attention-deficit/hyperactivity disorder) at the time of the offense for which they were incarcerated.[50]

Stimulant intoxication is marked by multiple indicators (see table S-1), which often present during withdrawal, as well. In addition, the behavioral signs of intoxication, such as psychosis, hallucinations, and delusions, can be difficult to

Screening

Recommendation

ii) Type used (e.g., methamphetamine, amphetamine, cocaine).

iii) Route of administration (intranasal, intravenous, oral, inhalation).

B. Frequency of use (regular daily or binge pattern).

C. Duration of the current period of regular use.

D. Concurrent use of other substances.

E. Psychiatric signs and symptoms.

F. Comorbid psychiatric illness.

S-4. Stimulant withdrawal is not usually associated with medical complications. However, long-term use of stimulants and stimulant intoxication are risk factors for cardiac complications. Health care and custody staff should be alert to chest pain and other cardiac symptoms in patients with suspected stimulant withdrawal.

S-5. Health care staff should also monitor for signs and symptoms of opioid withdrawal because the supply of stimulant drugs is increasingly contaminated or mixed with opioids, and some individuals may not be aware that they have been using opioids.

Monitoring Patients During Withdrawal Management

Recommendation

S-6. Substance withdrawal is a risk factor for severe depression and suicidality. This risk is particularly high in patients undergoing stimulant withdrawal. Custody staff should be alert to signs and symptoms of depression in these patients.

Table S-3: Possible Indicators of Depression [55]

- Aches or pains, headaches, cramps, or digestive problems without a clear physical cause and/or that do not ease even with treatment.
- Appetite and/or weight changes.
- Decreased energy or fatigue.
- Difficulty concentrating, remembering, or making decisions.
- Difficulty sleeping, early-morning awakening, or oversleeping.
- Feelings of guilt, worthlessness, or helplessness.
- Feeling restless or having trouble sitting still.
- Feelings of hopelessness or pessimism.
- Irritability.
- Loss of interest or pleasure in activities.
- Moving or talking more slowly.
- Persistent sad, anxious, or “empty” mood.
- Thoughts of death or suicide, or suicide attempts.

Medications

Treatment for stimulant withdrawal typically consists of behavioral management strategies and, when necessary, medications for symptom relief. Patients may present with mental health symptoms such as depression, agitation, suicidality, or stimulant-induced psychosis requiring medical management. For more information on suicide and SUD withdrawal, see Suicide in General Guidance.

Behavioral strategies, including providing an environment with reduced stimulation with dimmed lights, less noise, and minimal exposure to other people, can often reduce or prevent agitation. Care should be taken to encourage calm interactions and avoid situations that may become confrontational during the acute withdrawal phase. This may be challenging in a jail environment but can provide significant benefits, including reduced violence. Of note, the use of physical restraints can pose specific health risks in patients experiencing stimulant withdrawal.

Recommendations

S-7. No medications have been proven effective for the treatment of withdrawal from stimulants. However, it may be appropriate to use medications to relieve symptoms (e.g., agitation, sleep disturbances) during the period of the withdrawal syndrome.

S-8. The use of medications should be limited to short-term treatment of withdrawal symptoms and to treat accurately and appropriately diagnosed comorbid conditions.

S-9. If psychosis (beyond mild paranoia) manifests during stimulant withdrawal, it should be treated.

A. Care should be managed in consultation with a qualified mental health care professional.

B. Antipsychotic medication may be required.

C. If psychosis persists or is severe, an immediate assessment by a physician experienced in differential diagnosis and management of acute changes in mental status is indicated, and general psychosis management and treatment principles should be applied.

S-10. Patients experiencing agitation that does not immediately respond to behavioral management strategies (e.g., minimizing environmental stimulation [e.g., noise, bright lights, crowding]; speaking calmly to the patient; taking time to listen to and address concerns where appropriate and feasible) should receive an immediate medical evaluation.

A. Medications for sedation (typically benzodiazepines) may be necessary.

B. Patients whose agitation is not adequately treated with oral medication should be transferred to a hospital setting.

C. If medications are not available onsite to manage significant agitation, custody staff should transfer the patient to a medical setting where medications can be provided.

S-11. Benzodiazepines should be avoided unless required for concomitant alcohol or sedative detoxification, or severe agitation as discussed in the preceding recommendation.

S-12. While depression is common during stimulant withdrawal, it is often secondary to withdrawal and may not

Level of Care

Withdrawal from stimulants that is not complicated by other conditions can typically be safely managed in jails. However, ongoing or worsening severe depression and other psychiatric complications, as well as acute medical conditions (e.g., cardiac issues), may require transfer to a higher level of care.

Recommendations

S-13. As discussed in General Guidance, patients who present with severe psychiatric symptoms including hallucinations, delusions, paranoia, and delirium, should be immediately transferred to a higher level of care unless a physician experienced in differential diagnosis of acute changes in mental status is immediately available for clinical assessment and stabilization.

A. If feasible, medications may be appropriate to stabilize the patient concurrent with the call for transport.

S-14. Transfer to a higher level of care should occur when:

A. The patient displays significant psychiatric complications, and a mental health assessment cannot be provided

S-18.

Appendixes

Appendix A: Development Team

Appendix B: Full Methodology

Appendix C: Preferred Terminology

Appendix D: ASAM's Principles of Drug Testing During Withdrawal Management

Appendix E: Monitoring Patients During Withdrawal Management

Appendix F: Withdrawal Pharmacotherapy

Appendix G: Indicators of Dehydration

Appendix H: Expert Committee Members' Current Industry Relationships

Appendix I: Field Reviewers

Appendix J: Field Reviewers' Current Industry Relationships

Appendix K: Glossary

Appendix L: Acronym Glossary

Appendix M: Resources

Appendix A: Development Team

Expert Committee

Clinical Experts

Jeffrey Alvarez, M.D., CCHP
Chief Medical Officer, NaphCare

Andrew F. Angelino, M.D.
Associate Professor of Psychiatry and Behavioral Sciences, Johns Hopkins University of Medicine

Kevin A. Fiscella, M.D., M.P.H.
Addiction Medicine Physician, Jordan Health

Miriam S. Komaromy, M.D.
Medical Director, Grayken Center for Addiction at Boston Medical Center

Shannon Robinson, M.D.
Principal, Health Management Associates (HMA)

Sandra Springer, M.D.
Associate Professor of Medicine in the Department of Internal Medicine,
Section of Infectious Diseases Yale School of Medicine

Marc F. Stern, M.D., M.P.H.
Affiliate Assistant Professor, School of Public Health, University of Washington
Faculty, School of Public Health University at Albany
Chief Medical Advisor for the National Sheriffs' Association (NSA), American Jail Association (AJA),
and Washington Association of Sheriffs and Police Chiefs

Donna Strugar-Fritsch, B.S.N., M.P.A., CCHP
Correctional Health Care Consultant

Experts in Jail Administration

Oscar Aviles, CPM, CJM, CCE, CCHP
Assistant County Administrator, Hudson County (New Jersey)

Jeff Clark, M.J.A.
Undersheriff, Washoe County Sheriff's Office

Carrie Hill, Esq.
Executive Director, Massachusetts Sheriffs' Association
Chief Jail Advisor, NSA

Geoff Stobart
Chief Deputy of Research, Development, and Major Projects for the Franklin County Sheriff's Office (Columbus, Ohio)

Federal Project Champions

Stephen Amos, M.S.
National Institute of Corrections (NIC) Technical Lead
Chief, Jails Division
NIC, U.S. Department of Justice (DOJ)

Margaret Chapman, M.A.
Bureau of Justice Assistance Project Manager
Policy Advisor
BJA, DOJ

Ruby Qazilbash, M.P.A.
BJA Assistance Technical Lead
Deputy Director
BJA, DOJ

Technical Support Team

Maureen Boyle, B.A., Ph.D.
Executive Scientific Lead
Chief Quality and Science Officer, ASAM

Linda Frazier, B.S., M.A., RN, MCHES
Clinical Liaison
Director, Addictions Initiatives, Advocates for Human Potential, Inc. (AHP)

Brent Gibson, M.D., M.P.H., CCHP-P
Former Chief Health Officer
National Commission on Correctional Health Care (NCCHC)

Wendy Habert, M.B.A., PHR, CCHP
Director of Accreditation, NCCHC

Laura Lutgen-Nieves, Ph.D.
Literature Review Author
Assistant Professor, Department of Criminal Justice, University of Southern Indiana

Appendix B: Full Methodology

Overview of Approach

These guidelines were developed using a modified RAND/UCLA Appropriateness Method (RAM). The RAM process is a deliberate approach encompassing review of existing clinical guidelines and standards, literature reviews, appropriateness and feasibility ratings, stakeholder comment and reconciliation, and document development. The process typically combines scientific evidence and clinical knowledge to determine the appropriateness of a set of clinical procedures. This process was modified for the development of these guidelines to incorporate the input of jail administration experts and others with legal expertise regarding implementation of guidelines and procedures in jail settings.

RAM is particularly appropriate for the matter at hand for two reasons. First, there are few randomized clinical trials

The six guidelines listed below represented high quality clinical guidance addressing the broad clinical issues related to management of withdrawal from alcohol, opioids, sedative/hypnotics, and stimulants, and addressing management of withdrawal in closed settings flagged for EC members.

- *The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder, 2020 Focused Update* (2020)
- *Detoxification and Substance Abuse Treatment: Treatment Improvement Protocol (TIP) 45* (2015), from SAMHSA
- *Medically Supervised Withdrawal for Inmates with Substance Use Disorders* (2020), from the Federal Bureau of Prisons
- *The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management* (2020)
- *Clinical Guidelines for Withdrawal Management and Treatment of Drug Dependence in Closed Settings* (2009), from the World Health Organization
- *Models of Intervention and Care for Psychostimulant Users* (2004), from the Australian Government Department of Health and Ageing

These clinical documents were each developed using standard methodologies incorporating structured literature reviews and formal expert consensus development processes. They were used as the foundation for the development of draft recommendation statements (discussed below).

The EC was provided with the literature review summary, tables of key abstracted information from the scientific literature, and the guidelines identified in the gray literature search. This information was used by the EC to inform their discussion and judgments in Task 2.

Task 2: Identi cation of Draft Statements and Appropriateness/ Feasibility Rating

The next step in the RAM was to develop a set of draft recommendation statements, which were derived from the guideline analysis and informed by the literature review and EC's collective expertise. Each member of the EC reviewed the guidelines and literature review. The EC worked with the technical support team to draft recommendation statements. The EC then independently rated the 744 draft recommendation statements on a 9-point scale. The clinical experts on the committee rated the statements for appropriateness, and the jail administration experts rated the statements for feasibility within a jail setting.

A statement was considered appropriate if the expected health benefit (e.g., increased life expectancy, relief of pain, reduction in anxiety, improved functional capacity) exceeded the expected negative consequences (e.g., mortality,

Appendix C: Preferred Terminology

Use of person-first language and preferred terminology, which is reflected throughout these guidelines, reduces stigma, negative bias, and the perpetuation of stereotypes when speaking about or to individuals experiencing withdrawal or with substance use disorder (SUD).

Use …	Instead of …	Because …
• Individuals who are detained in jail • Individuals who enter jail • Person who has criminal justice system involvement • Person who was formerly incarcerated	• Inmate • Criminal • Offender • Justice-involved person • Felon	• Recognizes that people are not their crimes or criminal history. • Advances language that is accurate and not prejudicial.
• Substance use disorder • Substance use	• Substance abuse • Drug habit	• Avoids negative connotations and the undermining of SUD as a serious health condition.
• Person with a substance (opioid, alcohol, stimulant) use disorder • Individuals who exhibit signs of withdrawal or withdrawal symptoms • Individuals who report using substances (alcohol, opioids, stimulants) • Pregnant person with an opioid use disorder	• Addict • Drug or substance user • User • Dope addict • Alcoholic • Drug addict • Pregnant opiate addict	• Reflects an accurate, science-based understanding that the person has a medical condition. • Recognizes that the person has a disorder, rather than "is the disorder." • Avoids negative associations, punitive attitudes, individual blame, judgment, and misinformation about SUD. • Reinforces use of the same terminology as with other medical conditions. • Encompasses the recovery process of improved health versus the implication that the person was "dirty" before recovery.
• Individuals who screen negative/positive (e.g., withdrawal, suicide risk) • Positive/negative screen result	• Dirty/clean screen • Dirty results • Failing drug screen/test	
• Person in recovery • Person in long-term recovery • Person who has maintained recovery	• Former addict • Former drug user • Former alcoholic • Person who "stayed clean"	
• Return to use • Experienced a recurrence • Resumed drinking or using drugs	• Relapsed • Slipped • Chronic relapser	• Advances language that is medically accurate and non- stigmatizing.

Compiled from the Bureau of Justice Assistance, the National Institute on Drug Abuse, the Justice Community Opioid Innovation Network, and the National Council for Mental Wellbeing.

Appendix D: ASAM's Principles of Drug Testing During Withdrawal Management

1. Drug tests can be used to help inform clinical decisionmaking for patients with substance use disorder (SUD) or at risk for substance withdrawal.

2. Drug tests can neither diagnose SUD nor rule out SUD.

3. Drug test r

6. Definitive testing should be used when the results inform clinical decisions with major clinical or non-clinical implications for the patient (e.g., changes in medications, changes in legal status).

7. Drug test results should be interpreted by a provider whose scope of practice includes ordering and interpreting drug test results, who will consider the limitations of the specific test used.

8. Discrepancies between the patient self-report and the drug tests should be discussed with the patient.

9. Providers should keep test results ordered by health care staff confidential to the extent permitted by law.

10. Providers should be aware of the adverse legal and social consequences of detecting substance use in pregnant patients. The patient should be made aware of local/state reporting requirements and provide consent before tests are conducted.

For more information, go to ASAM's *Appropriate Use of Drug Testing in Clinical Addiction Medicine*.

Appendix E: Monitoring Patients During Withdrawal Management

WARNING: Supply patterns across the country show that drugs used in the community are often mixed with illicitly manufactured fentanyl and other drugs, such as xylazine. The most frequent monitoring noted is the minimum expectation, but providers may need to increase monitoring, based on their best clinical judgment and knowledge of local drug-use trends.

Substance	Minimum Frequency and Use of Withdrawal Symptom Assessment Scale*
Alcohol	Re-assessment using the Clinical Institute Withdrawal Assessment for Alcohol Scale, Revised (CIWA-Ar) at least every 8 hours during alcohol withdrawal management until the CIWA-Ar score remains below 10 for 24 hours. • If the CIWA-Ar is ≥ 19, repeat the CIWA-Ar at least every 6 hours during alcohol withdrawal management until the score falls below 19, and then continue monitoring with the CIWA-Ar at least every 8 hours until the score remains below 10 for 24 hours.
Sedatives	• Daily clinical assessment by a qualified health care professional for at least the first week or as condition

Appendix F: Withdrawal Pharmacotherapy

No medications have been approved by the FDA to treat stimulant withdrawal; therefore, that section describes behavioral management strategies as first-line treatment. Stocking medications for symptom relief is also recommended.

Table 2: Withdrawal Pharmacotherapy

Substance	Recommended Pharmacotherapy	For More Information
Alcohol	Benzodiazepines, long-acting preferred (i.e., diazepam or chlordiazepoxide), except for older adults or individuals with impaired liver function	*The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management*
Opioid	Methadone or buprenorphine	*The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update*
Sedatives	Long-acting benzodiazepine (e.g., clonazepam)	*Detoxification and Substance Abuse Treatment*, Treatment Improvement Protocol (TIP) Series, No. 45

Appendix I: Field Reviewers

American Academy of Addiction Psychiatry (AAAP)
Elie G. Aoun, M.D., MRO, FAPA
General, Addictions and Forensic Psychiatrist
Assistant Professor of Clinical Psychiatry, Columbia University, Division of Law, Ethics, and Psychiatry
Co-chair, AAAP Committee on Law and Addiction

Laurence M. Westreich, M.D.
Addiction and Forensic Psychiatry
Co-chair, AAAP Committee on Law and Addiction
Associate Professor of Clinical Psychiatry
Department of Psychiatry, New York University School of Medicine

American College of Academic Addiction Medicine (ACAAM)
Marsha J. Wunsch, M.D., FAAP DFASAM
Past President, ACAAM, Addiction Medicine Consultant
Alameda County Santa Rita Jail

American College of Medical Toxicology (ACMT)
Joseph Carpenter, M.D.
ACMT Addiction Medicine Committee, Communication Subcommittee Chair
Assistant Professor of Emergency Medicine, Emory University School of Medicine

Leslie R. Dye, M.D., FACMT, FASAM, FACCT
ACMT Addiction Toxicology Committee Chair and Past-President
Addiction Medicine Section Editor and former Editor-in-Chief, *Journal of Medical Toxicology*

Ashley Haynes, M.D.
ACMT Addiction Medicine Committee, Practice Subcommittee Chair
Program Physician, Wichita Comprehensive Treatment Center

American College of Obstetricians and Gynecologists (ACOG)
Carolyn Sufrin, M.D., Ph.D.
Associate Professor of Gynecology and Obstetrics, Johns Hopkins School of Medicine

Tricia Wright, M.D., M.S., FACOG DFASAM
Professor of Clinical Medicine, University of California San Francisco

Christopher M. Zahn, M.D.
Chief, Clinical Practice and Health Equity and Quality, ACOG

American Correctional Association (ACA)
Jeffrey Washington
Deputy Executive Director, ACA

American Jail Association (AJA)
Diana L. Knapp
Director, Jackson County Sheriff's Office

Henry Reyes, M.A.M., M.P.A., CJM, CCE, NCCE

External Reviewer	Representing	Salary	Consultant	Expert Witness	Ownership, Partnership, or Principal	Institution, Organization, or Other Financial Benefit
Darren Sieger, M.S., CJM	AJA	Broward Sheriff's Office, Department of Detention	None	None	None	None
Carolyn Sufrin, M.D., Ph.D.	ACOG	Johns Hopkins School of Medicine	None	None	None	NCCHC Board of Directors
Mishka Terplan N 185. 1 575.ID 386	EMC /P <</Lang (en-/P <<8/MCIDLan.671 /M	SAM0 1BT8.1l8/P	<</Lang dvisorycial)TjET	EMC /P <</Lang	(en-US)/MCI8hns Hopkins Scho	

Appendix K: Glossary

Definitions in this glossary were largely derived and adapted from evidence-based guidance documents and other reputable, field-accepted sources, as well as input from clinical and criminal justice experts.

Alcohol use disorder: A substance use disorder involving alcohol.[58]

Alcohol withdrawal delirium (formerly known as delirium tremens): A severe manifestation of alcohol withdrawal involving sudden and severe mental or nervous system changes. Clinical features include hallucinations, acute disorientation and confusion, autonomic instability, agitation, and paranoia.[59, 60]

Alpha-2 adrenergic agonists: A class of drugs (FDA-approved lofexidine and off-label clonidine) that activate alpha-2 adrenoceptors. These medications are often used to treat anxiety and hypertension and may also be used to help manage withdrawal symptoms when clinically appropriate.[61, 62]

Anxiolytics: A class of medications that help reduce the symptoms of anxiety.[63]

Appears unwell:

Complicated withdrawal: Severe withdrawal symptoms involving seizures, hallucinations, delirium, or psychosis, which require more frequent monitoring and may require hospital-level care.

Cravings: Desire to use substances or engage in addictive behaviors, experienced as a physical or emotional need for reward and/or relief.[73]

Custodial facility: Refers to confines such as jail, detention, holding, or lockup facilities.

Custody staff:

Licensed prescriber: A nurse practitioner, physician assistant, physician, or other health care provider licensed to prescribe medications. Note that state laws may vary with regard to prescribing authorities.

Methadone treatment provider: Authorized provider of methadone (e.g., opioid treatment programs are certified by SAMHSA to treat patients with opioid use disorder using methadone).

Micro-dosing: Administration of buprenorphine–naloxone in a small initial dose with incremental increases to both dose and frequency over time.[87]

Misuse: Misuse of prescription drugs means taking a medication in a manner or dose other than prescribed; taking someone else's prescription, even if for a legitimate medical complaint such as pain; or taking a medication to feel euphoria (i.e., to "get high").[88]

Monitoring: Regular and active surveillance to detect changes in physical or mental status that may indicate health problems, which facilitates appropriate interventions and ensures patient safety.[89, 90]

Narcotic treatment program: The Controlled Substances Act and U.S. Drug Enforcement Administration (DEA) regulations refer to opioid treatment programs as narcotic treatment programs, which are registered with the DEA.[91]

Older adult: Individuals who are 55 years and older.[92]

Opioid: Any psychoactive chemical resembling morphine in pharmacological effects, including opiates and synthetic/semisynthetic agents.[93]

Opioid agonist: A medication that occupies and activates opioid receptors in the body.[94]

Opioid antagonist: A medication that occupies opioid receptors in the body but does not activate the receptors. This effectively blocks the receptor, preventing the brain from responding to other opioids. The result is that further use of opioids does not produce analgesia, euphoria, or intoxication.[95]

Opioid use disorder: A substance use disorder involving opioids.[96]

Opioid withdrawal syndrome: The wide range of symptoms occurring when individuals who regularly take opioids, either illicit or prescribed, stop or reduce their use.[97]

Patient: As used in this document, a person whose substance withdrawal or SUD is being treated.[98]

Patient navigator: A person whose role is to help individuals who are transitioning to new circumstances understand system processes and how to effectively navigate systems to obtain services needed and access resources; for purposes of this document, this transition is from jail to the community. Often, patient navigators have lived experience with a substance use disorder and/or involvement with the criminal justice system.[99]

Pharmacokinetics: The study of how pharmacological agents are processed within a biological system, including factors that influence the absorption, distribution, metabolism, and elimination of a substance or its metabolic products.[100]

Pharmacotherapy: Therapy (medical treatment) using pharmaceutical drugs.[101]

Physician: A designated doctor of medicine or osteopathy who has the final authority at a given facility regarding clinical issues.

Physiological dependence: Adaptation of a body to a substance, requiring more of it to achieve a certain effect (tolerance) and eliciting substance-specific physical or mental symptoms if drug use is abruptly ceased (withdrawal).

Appendix L: Acronym Glossary

AAAP	American Academy of Addiction Psychiatry
ACA	American Correctional Association
ACAAM	American College of Academic Addiction Medicine
ACMT	American College of Medical Toxicology
ACOG	American College of Obstetricians and Gynecologists
AHP	Advocates for Human Potential, Inc.
AJA	American Jail Association
AOAAM	American Osteopathic Academy of Addiction Medicine
ASAM	American Society of Addiction Medicine
AUD	Alcohol use disorder
BAC	Blood alcohol concentration
BJA	Bureau of Justice Assistance
BrAC	Breath alcohol concentration
CAPTA	Child Abuse Prevention and Treatment Act
CIWA-Ar	Clinical Institute Withdrawal Assessment for Alcohol Scale, Revised
COCHS	Community Oriented Correctional Health Services
COPS	Office of Community Oriented Policing Services
COSSUP	BJA's Comprehensive Opioid, Stimulant, and Substance Use Program
COWS	Clinical Opiate Withdrawal Score
CPR	Cardiopulmonary resuscitation
CRD	Civil Rights Division
DEA	U.S. Drug Enforcement Administration
EC	Expert committee
EMS	Emergency medical services
FASD	Fetal alcohol spectrum disorder
FDA	U.S. Food and Drug Administration
GHB	Gamma-hydroxybutyric acid
HMA	Health Management Associates

IV	Intravenous
MCSA	Major County Sheriffs of America
MOU	Memorandum of understanding
NAATP	National Association of Addiction Treatment Providers
NACo	National Association of Counties
NCCHC	National Commission on Correctional Health Care
NIAAA	National Institute on Alcohol Abuse and Alcoholism
NIDA	National Institute on Drug Abuse
NLM	National Library of Medicine
NPG	National Practice Guideline
NSA	National Sheriffs' Association
NSAID	Nonsteroidal anti-inflammatory medication (drug)
OTP	Opioid treatment program
OUD	Opioid use disorder
PDMP	Prescription Drug Monitoring Program
RAM	RAND/UCLA Appropriateness Method
RCT	Randomized clinical trial
SAMHSA	Substance Abuse Mental Health Services Administration
SRLEEA	Small & Rural Law Enforcement Executives Association
SUD	Substance use disorder
TIP	Treatment Improvement Protocol
VA	U.S. Department of Veterans Affairs

Appendix M: Resources

To navigate to various sections of this resource list, click on the corresponding hyperlink.

Clinical Practice Studies
Legislation, Rules, and Regulations
Pregnancy and Postpartum
Substances
Treatment and Medication
Reentry
Screening and Assessment
Suicide
Telehealth

Clinical Practice Studies

"Microinduction of Buprenorphine/Naloxone: A Review of the Literature"—review of existing literature to help clinicians better understand the approaches to micro-dosing of buprenorphine in various clinical settings and populations. (*American Journal on Addictions*)

"Review Article: Rapid Review of the Emergency Department-initiated Buprenorphine for Opioid Use Disorder"—investigation of the effectiveness of initiating buprenorphine in the emergency department setting. (*Emergency Medicine Australasia*)

Treatment for Pregnant People with Opioid Use Disorder in Jail—study focusing on jails, recognizing that these facilities

Managing Substance Withdrawal in Jails: A Legal Brief—the scope of challenge, overview of key legislation and court cases related to substance withdrawal, and steps to create a comprehensive response for people with SUD. (Bureau of Justice Assistance [BJA], National Institute of Corrections)

Pregnancy and Postpartum

Clinical Guidance for Treating Pregnant and Parenting Women with Opioid Use Disorder and Their Infants—comprehensive guidance for optimal management of pregnant and parenting people with OUD and their infants. (Substance Abuse and Mental Health Services Administration [SAMHSA])

Opioid Use and Opioid Use Disorder in Pregnancy—recommendations for identification of and treatment for pregnant persons with SUD. (The American College of Obstetricians and Gynecologists [ACOG])

Reproductive Health Care for Incarcerated Pregnant, Postpartum, and Nonpregnant Individuals—how obstetricians-gynecologists and other practitioners can support efforts to improve health care for pregnant, postpartum, and non-pregnant persons who are incarcerated. (ACOG)

Substance Abuse Reporting and Pregnancy: Role of the Obstetrician-Gynecologist—discussion of reporting requirements related to SUD that an obstetrician-gynecologist should be aware of within their state. (ACOG)

Substance Use and Pregnancy – Part 1: Current State Policies on Mandatory Reporting and Implementing Plans of Self Care to Support Pregnant Persons with Substance Use Disorders—examination of state-level policies supporting or inhibiting pregnant and postpartum persons' access to long-term recovery from OUD. (BJA's Comprehensive Opioid, Stimulant, and Substance Use Program [COSSUP])

Substance Use During Pregnancy and Family Care Plans—fact sheet on substance use during pregnancy and family planning. (Legislative Analysis and Public Policy Association with funding from the Office of National Drug Control Policy)

Substances

Barbiturates—fact sheet on barbiturates. (Drug Enforcement Administration [DEA])

Benzodiazepines—description of benzodiazepines, including its licit and illicit uses. (DEA)

Chlordiazepoxide—description of chlordiazepoxide and its uses. (National Library of Medicine's [NLM's] MedlinePlus)

Clonazepam—description of clonazepam and its uses. (NLM's Medline Plus)

Diazepam—description of diazepam and its uses. (NLM's MedlinePlus)

Drug Fact Sheets—descriptions of various medications, as well as illicit drugs. (DEA)

Facts About Fentanyl—information about fentanyl. (DEA)

Gamma Hydroxybutyric Acid (GHB)—description of GHB, including its licit and illicit uses. (DEA)

Substance Use – Prescription Drugs—common medications that are misused and a list of "street names." (NLM's MedlinePlus)

Opioid Treatment Program Directory—listing of opioid treatment programs by state. (SAMHSA)

Opioid Use Disorder Treatment in Correctional Settings—position statement on patient health records upon release. (NCCHC)

Performance Measures for Medication-assisted Treatment in Correctional Settings: A Framework for Implementation—framework for jail and prison administrators, program managers, medical staff in correctional settings, and reentry staff to monitor MAT in correctional settings. (Legislative Analysis and Public Policy Association)

A Primer for Implementation of Overdose Education and Naloxone Distribution in Jails and Prisons—strategies for developing, coordinating, monitoring, and evaluating jail and prison-based programs, as well as lessons learned from two studies. (RTI International with funding from NIDA)

Standards for Opioid Treatment Programs—requirements for corrections-based opioid treatment programs seeking accreditation. (NCCHC)

Substance Abuse Treatment: Addressing the Specific Needs of Women, TIP 51—guidance for providers in offering treatment to women with SUD. (SAMHSA)

Treating Substance Use Disorder in Older Adults, TIP 26—guidance for providers and others to better understand how to identify, manage, and prevent substance misuse among older adults. (SAMHSA)

Treatment for Stimulant Use Disorders, TIP 33—recommendations on treatment approaches, strategies for planning and initiating treatment, as well as how to maximize treatment engagement and retention, and strategies for initiating and maintaining abstinence. (SAMHSA)

Use of Medication-assisted Treatment for Opioid Use Disorder in Criminal Justice Settings—use of MAT for OUD in jails and prisons and during the reentry process, providing an overview of policies and evidence-based practices reducing the risk of overdose and relapse. (SAMHSA)

Reentry

Guidelines for Successful Transition of People with Mental or Substance Use Disorders from Jail and Prison: Implementation Guide—behavioral health, correctional, and community stakeholders with implementation examples for transitioning people with mental disorders or SUD from correctional settings into the community. (SAMHSA)

National Reentry Resource Center—primary source of reentry information and guidance. (BJA)

Optimizing Insurance Coverage for Individuals Postrelease—position statement on insurance coverage upon release. (NCCHC)

Principles of Community-based Behavioral Health Services for Justice-involved Individuals: A Research-based Guide—guidance for community-based behavioral health providers in their clinical and case management practice. (SAMHSA)

Sharing of Patient Health Records Upon Release from Incarceration—position statement on patient health records upon release. (NCCHC)

Screening and Assessment

CIWA-Ar (Clinical Institute Withdrawal Assessment of Alcohol Scale, Revised)—commonly used withdrawal assessment tool for alcohol. (ASAM)

COWS

Telehealth

Telehealth Implementation Support Tool—an online support tool intended to be completed in approximately 20 minutes by a jail administrator, who may need input from other stakeholders. (COSSUP)

"Telemedicine," from the COVID-19 Information web page—guidance on how telemedicine can be used, including the DEA policy on the use of telephone evaluations to initiate buprenorphine. (DEA)

Using Telehealth for Behavioral Health in the Criminal Justice System—a brief on telehealth services and its benefits to the criminal justice system as an innovative strategy for intervention and treatment of OUD. (COSSUP)

Endnotes

[1]The White House, Executive Of ce of the President, Of ce of National Drug Control Policy, 2022, *National Drug Control Strategy*: 98, retrieved April 12, 2023 from https://www.whitehouse.gov/wp-content/uploads/2022/04/National-Drug-Control-2022Strategy.pdf .

[2]Bronson, Jennifer, Jessica Stroop, Stephanie Zimmer, and Marcus Berzofsky, June 2017, *Drug Use, Dependence, and Abuse Among State Prisoners and Jail Confined Persons, 2007–2009*, Washington, DC: U.S. Department of Justice, Bureau of Justice Statistics: 3, NCJ 250546, retrieved February 9, 2022 from https://bjs.ojp.gov/content/pub/pdf/dudaspji0709.pdf .

[3]Fiscella, Kevin, Margaret Noonan, Susan H. Leonard, Subrina Farah, Mechelle Sanders, Sarah E. Wakeman, and Jukka Savolainen, 2020, "Drug- and Alcohol-Associated Deaths in U.S. Jails," *Journal of Correctional Health Care 26*(2): 183–193, retrieved March 19, 2021 from http://doi. org/10.1177/1078345820917356 .

[4]E. Ann Carson, December 2021, Mortality in Local Jails, 2000–2019 – Statistical Tables, Washington, DC: U.S.

[11]Substance Abuse and Mental Health Services Administration, 2017, Peers Supporting Recovery from Substance Use Disorders, Rockville, MD: U.S. Department of Health and Human Services, retrieved February 3, 2023 from https://www.samhsa.gov/sites/default/ les/programs_campaigns/brss_tacs/peers-supporting-recovery-substance-use-disorders-2017.pdf .

[12]S m 25.3701 Tm (12)Tj ET 0 k tmegen SeO8 (, 20cians, r)1Gynse-logj Es' k mmitteen neO8 (, 20c PrBDCd F in e-llaboT EMC w

[37]American Society of Addiction, "ASAM's Treatment of Opioid Use Disorder with Buprenorphine: Clinical Considerations for Treatment of Individuals with OUD Using High Potency Synthetic Opioids," Journal of Addiction Medicine , submitted March 2023.

[38]See note 32 above, American Society of Addiction Medicine, The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update , 9.

[39]Code of Federal Regulations, 2023, Administering or dispensing of narcotic drugs , Title 21, Chapter ll, Part 1306.07(b), retrieved February 7, 2023 from https://www.ecfr.gov/current/title-21/chapter-II/part-1306/subject-group-ECFR1eb5bb3a23fddd0/section-1306.07 .

[40]Substance Abuse and Mental Health Services Administration, n.d., Certi cation of Opioid Treatment Programs (OTPs) , retrieved March 31, 2023 from https://www.samhsa.gov/medications-substance-use-disorders/become-accredited-opioid-treatment-program .

[41]Substance Abuse and Mental Health Services Administration, 2021, Medications for Opioid Use Disorder , Treatment Improvement Protocol (TIP) 63, Rockville, MD: U.S. Department of Health and Human Services: Chapter 3B, Publication No. PEP21-02-01-002, retrieved March 8, 2022 from https://store.samhsa.gov/sites/default/ les/SAMHSA_Digital_Download/PEP21-02-01-002.pdf .

[42]DailyMed, 2007, LABEL: Methadone- Methadone Hydrochloride Injection, Solution , Bethesda, MD: U.S. National Library of Medicine, retrieved March 8, 2022 from https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=9bbe219f-3bf1-44f4-a709-cd108c4fa9ef .

[43]Civil Rights Division, 2022, The Americans with Disabilities Act and the Opioid Crisis: Combating Discrimination Against People in Treatment or Recovery , Washington, DC: U.S. Department of Justice, retrieved June 21, 2022 from https://archive.ada.gov/opioid_guidance.pdf .

[44]See note 32 above, American Society of Addiction Medicine, The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update , 61.

[45]Fox, Aaron D., Jeronimo Maradiaga, Linda Weiss, Jennifer Sanchez, Joanna L. Starrels, and Chinazo O. Cunningham, 2015, " Release from Incarceration, Relapse to Opioid Use and the Potential for Buprenorphine Maintenance Treatment: A Qualitative Study of the Perceptions of Former Inmates with Opioid Use Disorder ," Addiction Science & Clinical Practice, 10:2, retrieved April 4, 2023 from https://ascpjournal.biomedcentral.com/articles/10.1186/s13722-014-0023-0 .

[46]See note 32 above, American Society of Addiction Medicine, The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update , 49.

[47]See note 12 above, American College of Obstetricians and Gynecologists' Committee on Obstetric Practice in collaboration with Maria Macola, Ann Borders, and the American Society of Addiction Medicine Mishka Terplan, Opioid Use and Opioid Use Disorder in Pregnancy .

[48]Ibid.

[49]Malta, Monica, Thepikaa Varatharajan, Cayley Russell, Michelle Pang, Sarah Bonato, and Benedikt Fischer, 2019, "

[50]Maruschak, Laura M., Jennifer Bronson, and Mariel Alper, July 2021, Survey of Prison Inmates, 2016: Alcohol and Drug Use and Treatment Reported by Prisoners, Statistical Tables, Washington, D.C.: U.S. Department of Justice, Bureau of Justice Statistics: 2, NCJ 252641, retrieved March 18, 2022 from https://bjs.ojp.gov/sites/g/ les/xyckuh236/ les/media/ document/adutrpspi16st.pdf .

[51]Mahoney, James J., Ari D. Kalechstein, Richard De La Garza, and Thomas F. Newton, 2008, "Presence and Persistence

[62]MedlinePlus, n.d., Clonidine, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 24, 2022 from https://medlineplus.gov/druginfo/meds/a682243.html.

[63]National Institute of Mental Health, n.d., Mental Health Medications, Bethesda, MD: National Institutes of Health, U.S. Department of Health and Human Services, retrieved March 18, 2022 from https://www.nimh.nih.gov/health/topics/mental-health-medications.

[64]MedlinePlus, n.d., Asymptomatic, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 18, 2022 from https://medlineplus.gov/ency/article/002217.htm.

[65]MedlinePlus, n.d., Movement - Uncoordinated, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 18, 2022 from https://medlineplus.gov/ency/article/003198.htm.

[66]MedlinePlus, n.d., Autonomic Nervous System Disorders, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 18, 2022 from https://medlineplus.gov/autonomicnervoussystemdisorders.html.

[67]Skibiski, Jeanie, and Sara Abdijadid, 2022, Barbiturates, Treasure Island, FL: StatPearls Publishing, retrieved March 10, 2022 from https://www.ncbi.nlm.nih.gov/books/NBK539731.

[68]Ibid; MedlinePlus, 2020, Phenobarbital, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 14, 2022 from https://medlineplus.gov/druginfo/meds/a682007.html#:~:text=Phenobarbital%20is%20used%20to%20control,to%20stop%20taking%20the%20medication.

[69]National Institute on Drug Abuse, 2022, November 7, Benzodiazepines and Opioids, Rockville, MD: National Institutes of Health, retrieved March 13, 2022 from https://nida.nih.gov/research-topics/opioids/benzodiazepines-opioids.

[70]Drug & Chemical Evaluation Section, December 2019, Benzodiazepines (Street Names: Benzos, Downers, Nerve Pills, Tranks), Spring eld, VA: Drug Enforcement Administration, retrieved March 10, 2022 from https://www.deadiversion.usdoj.gov/drug_chem_info/benzo.pdf.

[71]Substance Abuse and Mental Health Services Administration, Screening and Assessment of Co-Occurring Disorders in the Justice System, 2015, Rockville, MD: U.S. Department of Health and Human Services, HHS Publication No. PEP19-SCREEN-CODJS, retrieved March 22, 2022 from https://store.samhsa.gov/sites/default/ les/d7/priv/pep19-screen-codjs.pdf.

[72]National Institute on Drug Abuse, n.d., Comorbidity, Rockville, MD: National Institutes of Health, retrieved March 11, 2022 from https://nida.nih.gov/research-topics/comorbidity.

[73]Mee-Lee, D., G.D. Shulman, M.J. Fishman, D.R. Gastfried, and M.M. Miller, eds., 2013, The ASAM Criteria: Treatment Criteria for Addictive, Substance-related, and Co-occurring Conditions. 3rd ed. Carson City, NV: The Change Companies: 417.

[74]National Commission on Correctional Health Care, 2018, Jail and Prison Standards, Chicago, IL: National Commission on Correctional Health Care: 6.

[75]MedlinePlus, n.d., Delirium, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 18, 2022 from https://medlineplus.gov/delirium.html.

[76]MedlinePlus, n.d., Differential Diagnosis, Bethesda, MD: National Library of Medicine, National Institutes of Health, retrieved March 23, 2023 from https://medlineplus.gov/lab-tests/differential-diagnosis.

manual_Final.pdf.

[92]Greene, Meredith, Cyrus Ahalt, Irena Stijacic-Cenzer, Lia Metzger, and Brie Williams, 2018, " Older Adults in Jail: High Rates and Early Onset of Geriatric Conditions ," Health Justice 6:3, 2194–7899, retrieved from https://doi.org/10.1186/s40352-018-0062-9.

[93]See note 32 above, American Society of Addiction Medicine, The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update , 5.

[94]Ibid., 5.

[95]Ibid., 5.

[96]Ibid., 6.

[97]Ibid., 6.

[98]See note 73 above, Mee-Lee et al., The ASAM Criteria: Treatment Criteria for Addictive, Substance-related, and Co-occurring Conditions , 425.

[99]Substance Abuse and Mental Health Services Administration, August 2017, Peer Support Roles in Criminal Justice Settings, Rockville, MD: U.S. Department of Health and Human Services, retrieved April 4, 2023 from https://www.cdcr.ca.gov/ccjbh/wp-content/uploads/sites/172/2019/06/WebinarSupportingDocument_PeerRolesinCJSettings508.pdf .

[100]APA Dictionary of Psychology, n.d., Pharmacokinetics, Washington, DC: American Psychological Association, retrieved March 11, 2022 from https://dictionary.apa.org/pharmacokinetics .

[101]See note 16 above, American Society of Addiction Medicine, The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management , 4.

[102]National Institute on Drug Abuse, January 2018, National Institute on Drug Abuse (NIDA) Principles of Drug Addiction Treatment: A Research-Based Guide (Third Edition), Rockville, MD: National Institutes of Health: 26, retrieved on March 18, 2023 from https://nida.nih.gov/sites/default/ les/675-principles-of-drug-addiction-treatment-a-research-based-guide-third-edition.pdf .

[103]See note 32 above, American Society of Addiction Medicine, The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update , 6.

[104]APA Dictionary of Psychology, n.d., Psychosis, Washington, DC: American Psychological Association, retrieved on March 11, 2022 from https://dictionary.apa.org/psychosis .

[105]National Commission on Correctional Ad1isorP </Lang L(en-US)PsyD 5465 >>BD5dC10 0 0 10 132.18(en-US)Ps/GS1 gs /T1_0

[107]APA Dictionary of Psychology, n.d.,

[123]APA Dictionary of Psychology, n.d., Wernicke's Encephalopathy, Washington, DC: American Psychological Association, retrieved March 11, 2022 from https://dictionary.apa.org/wernickes-encephalopathy.

[124]See note 73 above, Mee-Lee et al., American Society of Addiction Medicine, The ASAM Criteria: Treatment Criteria for Addictive, Substance-related, and Co-occurring Conditions, 432.

[125]Ibid., 432.

[126]Naren Gunja, 2013, "The Clinical and Forensic Toxicology of Z-drugs," Journal of Medical Toxicology 9: 155–162, retrieved September 14, 2022 from https://doi.org/10.1007/s13181-013-0292-0.

[127]Schifano, Fabrizio, Stefania Chiappini,/doi.org/10.1007/s13181-013-02 >>BDC BT 0 0 0 1 k /GS1 gs 136.6 610.4401 Tm [(r)18

Bureau of Justice As sistance Information

BJA helps to make American communities safer by strengthening the nation's criminal justice system; its grants, training and technical assistance, and policy development services provide state, local, and tribal governments with the cutting-edge tools and best practices they need to reduce violent and drug-related crime, support law enforcement, and combat victimization.

- Emphasize local control.
- Build relationships in the eld.
- Provide training and technical assistance in support of efforts to prevent crime, drug abuse, and violence at the national, state, and local levels.
- Develop collaborations and partnerships.
- Promote capacity building through planning.
- Streamline the administration of grants.
- Increase training and technical assistance.
- Create accountability of projects.
- Encourage innovation.
- Communicate the value of justice efforts to decision makers at every level.

To learn more about BJA, visit

bja.ojp.gov

Facebook (www.facebook.com/DOJBJA)

Twitter (@DOJBJA).

BJA is a component of the Department of Justice's
Of ce of Justice Programs.