February 24, 2020
12:30-2:00pm ET

Hosted by SAMHSA's GAINS Center

Lisa Callahan, PhD
Senior Research Associate II

The views, opinions, and content expressed in this presentation and discussion do not necessarily reflect the views, opinions, or policies of the Center for Mental Health Services (CMHS), the Center for Substance Abuse Treatment (CSAT), the Substance Abuse and Mental Health Services Administration (SAMHSA), or the U.S. Department of Health and Human Services (DHHS).

/ v š Œ } µ] v P bã pb•x'p •f 374.3tr.G77>-a

- ‡Is Senior Scientific Consultant for the National Association of Drug Court Professionals (NADCP), and Senior Science & Policy Advisor for Alcohol Monitoring Systems (AMS).
- ‡Formerly served as Chief of Science, Law & Policy for NADCP; Director of Law & Ethics Research at the Treatment Research Institute (TRI); and Adjunct Associate Professor of Psychiatry at the University of Pennsylvania School of Medicine.
- ‡Is a lawyer and clinical psychologist who studies the effects of treatment courts and other rehabilitation programs for persons with substance use and mental health disorders in the justice system.
- ‡Published over 175 journal articles, monographs, books, and book chapters on the topics of correctional rehabilitation, forensic psychology, and treatment of substance use disorders.
- ‡Serves as Editor-in-Chief of the Journal for Advancing Justice, is on the editorial board of Criminal Justice & Behavior, and was previously Editor-in-Chief of the Drug Court Review.

/všŒ} µ]vP d} Ç[• WŒ • vš Œ•W ^š À ,

‡Is the Associate Commissioner for Courts and Criminal Justice for the New York State (NYS) Office of Alcoholism and Substance Abuse Services.

‡Served previously as the NYS Associate Commissioner for Treatment, overseeing treatment services in the state of New York, including direct oversight of 12 state-operated inpatient treatment programs.

‡Has been on the faculty of the National Drug Court Institute at NADCP since 2001.

‡Has presented both nationally and internationally on the topics of Addiction and Treatment.

/všŒ} µ]vP d} Ç[• WŒ • vš Œ•W ^Z µv

‡Is a district court judge of the 6th Judicial District of Minnesota, chambered in Duluth, MN.

‡Founded and presides over the South St. Louis County DWI Court in Minnesota, one of four National Center for DWI Courts Academy Courts in the nation.

‡Is a member of the National Judicial Opioid Task Force and serves on the Judicial Advisory Board for the Foundation for Advancing Alcohol Responsibility.

‡Is the founding and presiding judge of the Duluth Domestic Violence Restorative Circles Intervention in Duluth, MN.

‡Serves as Co-chair of the Minnesota Treatment Court Initiative.

‡

Senior Scientific Consultant

National Association of Drug Court Professionals

February 24, 2020



Etc. . .

Domains	Methadone	Buprenorphine	XR-Naltrexone
[illegible]	√	O	E I
Opioid cravings	√	O	√
Treatment entry & retention	√	O	√
[illegible]	O	O	O
Criminal recidivism	E I	E I	√
[illegible]	√	√	√
[illegible]	√	E I	√

(SAMHSA, 2019)

- √ at least two meta-analyses, systematic reviews or experimental trials in criminal justice populations
- O experimental trials in non-criminal justice settings or correlational studies in criminal justice populations

‡Obligation to learn the facts about MAT

‡Obtain medical consultation

‡No blanket prohibitions for entry or graduation

‡

W D W • ~ } v š [•

‡ Drug courts should require inquiries in their memoranda of understanding (MOUs) with providers.

/v(}Œu š]}v }µš Ç}µŒ •š š }(

- Is PDMP enrollment mandatory or permissive in your jurisdiction?
  - Mandatory enrollment for N=44 (83%) of prescribers, dispensers or both as of Jan, 2019
  - Enrollment conditions by state
- Must providers query the PDMP for new covered prescriptions or refills?
  - N=45 (85%) of states required inquiries as of Jan, 2019
- Data-sharing reciprocity with other jurisdictions.
  - Interstate data-sharing partners
- Data-sharing with law enforcement, community corrections and drug courts.
  - Approx. one-third of states explicitly authorize reports to drug courts
  - Law enforcement access methods to PDMP reports
  - Solicited and unsolicited reports to law enforcement agencies

‡National Alliance for Model State Drug Laws



https://namsdl.org/wp-content/uploads/Established-and-Operational-Prescription-Drug-Monitoring-Programs-PMPs-–-Map.pdf

- Methadone may only be prescribed or dispensed for the treatment of substance use disorders from a licensed opioid treatment program (OTP).
  - Take-home doses may be permitted after meeting legally specified requirements for treatment attendance, clinical stability, and abstinence from illicit substances.

(SAMHSA, 2015)

µšZ}Œ]Ì D d WŒ}À] Œ• ~ }v

- Buprenorphine (e.g., Suboxone, Subutex, Probuphine, Sublocade) may only be prescribed by a DATA-2000 waivered medical provider.
  - Physicians must complete an 8-hour training and can treat up to 100 patients in the first year and 275 patients thereafter.
  - Nurse practitioners and physician's assistants must complete a 24-hour training and can treat up to 100 patients.
  - Only about 5% of physicians have obtained DATA-2000 Waivers.
- Naltrexone (including Vivitrol) may be prescribed and dispensed by licensed medical providers such as physicians, physician assistants, and nurse practitioners.

(SAMHSA, 2019; National Drug Court Institute, 2013; SAMHSA, nd)

‡ SAMHSA Behavioral Health Treatment Services Locator

‡ SAMHSA Buprenorphine Practitioner Locator

‡ SAMHSA Opioid Treatment Program Directory

‡ American Academy of Addiction Psychiatry (AAAP)

‡ American Board of Addiction Medicine (ABAM)

‡ American Society of Addiction Medicine (ASAM)

Providers may also be found through single-state agencies for substance use services, local colleges & medical schools, or county or state boards of health.

‡ SAMHSA Buprenorphine Training for Physicians

‡ SAMHSA Buprenorphine Waiver Management

‡ American Association of Nurse Practitioners (AANP)

‡ ASAM eLearning

‡ Providers Clinical Support System (PCSS)

‡ All participants should be screened routinely for symptoms of opioid use disorder, including current and past opioid use, withdrawal symptoms, cravings, overdose history, opioid treatment history, and past receipt of MAT.

‡ Any person screening positive should be referred to a licensed and trained medical provider for a follow-up diagnostic evaluation and determination of suitability for MAT.

(SAMHSA, 2015)

‡

- Texas Christian University (TCU) Drug Screen-5-Opioid Supplement
- Rapid Opioid Dependence Screen (RODS)
- Severity of Opioid Dependence Questionnaire (SODQ)

‡

- Clinical Institute Narcotic Assessment (CINA) Scale for Withdrawal Symptoms
- Clinical Opiate Withdrawal Scale (COWS)
- Subjective Opiate Withdrawal Scale (SOWS)

‡

- Brief Substance Craving Scale (BSCS)
- Heroin Craving Questionnaire (HCQ)
- Opioid Craving Scale (OCS)

‡

- Current Opioid Misuse Measure (COMM)
- _____

‡ Narcotics Anonymous (NA, 2016) policy is accepting of MAT, including agonists (methadone and buprenorphine); however, many groups disapprove of MAT or discourage entrance for those receiving it.

‡ In a recent study, 58% of drug court participants reported experiencing or witnessing disapproval of MAT from peer-

W Œ ^ µ ‰ ‰ } Œ š ' Œ } µ ‰ • ~ } v š [

‡ Preparing participants for possible disapproval and discussing how and whether to share their receipt of MAT increases group satisfaction and tenure (Galanter, 2018; Krawczyk et al., 2018; Suzuki & Dodds, 2016; White et al., 2013).

‡ Peer-support groups accepting of MAT can be found at Medication-Assisted Recovery Anonymous (MARA).

W Ç]vP (}Œ D d ~ }vš[•

‡ Drug court participants and people on probation are fully eligible.

- ƒ As of 2/2018, 36 states and territories covered methadone, 51 covered buprenorphine, 49 covered naltrexone, and over 50% increased coverage for naloxone (Narcan).
- ƒ There is discretion to cover “rehabilitative services”, broadly defined to include substance use counseling, peer support specialists, supportive housing, and vocational services.
- ƒ There is discretion to cover benefits assistants, case managers, and data exchanges.

(SAMHSA, 2019)

‡ MAT providers and criminal justice professionals meeting regularly for at least 12 months to discuss mutual concerns and resolve barriers increased actual referral intentions and rates (Friedmann et al, 2015).

‡ Equipping staff with naloxone kits, teaching them to assemble and use the applicators, and providing videotaped boosters increased actual uptake and distribution of naloxone kits (Anthony

‡ Use terms associated with greater empathy and optimism. (e.g., “recovery”, “recurrence of use”, “suffering from . . .”)

‡ Describe previously unsuccessful efforts at recovery without disheartened reactions (“devastated”).

‡ Use vignettes describing successful efforts at recovery.

‡ Describe practical misuse-prevention strategies.

‡ Educate about the disease model of addiction, including brain imaging & genetics.

‡ Describe common environmental and social precipitants (e.g., trauma, peer influences, poor prescription practices, exposure to advertising).

‡ Describe medical and psychiatric disorders that frequently co-occur with substance use disorders (e.g., chronic pain, depression, PTSD).

- Blanket prohibitions or routine denials of MAT violate the Americans with Disabilities Act (ADA), Rehabilitation Act, and possibly the 8th Amendment prohibition against "deliberate indifference" to medical needs of incarcerated people.
  - Pesce v. Coppinger, No. 18-11972-DJC (D. Mass. 2018); Smith v. Aroostook County, No. 1:18-cv-352-NT (D. Maine 2019), (([No. 19-1340 (1st Cir. 2019); DiPierro v. Hurwitz, No. 1:19-cv-10495-WGY (D. Mass.2019) (settlement agreement)
- At least one DOJ warning letter to a treatment court asserted that treating participants differently because they were receiving MAT violated the ADA.
- Sample letters to educate jail officials and others: https://lac.org

There are several common obstacles to implementing Medication-assisted Treatment (MAT) in Drug Courts, including:

- ‡Stigma
- ‡Logistics/Access
- ‡Cost
- ‡Diversion concerns

Stigma is the largest obstacle to MAT. The stigma is based on “philosophical” bias and erroneous beliefs about the medication (Wakeman and Rich, 2017).

‡ Many people have achieved Recovery without medication.
t

‡ If you have to take a medication, it’s not “real recovery.”
t

‡ You are just replacing one drug with another.
t E } U] š [• u] š] } v X

Œ ••]vP ŒŒ}v }µ• o] (• ~

‡ People on MAT are just zombies.

t

‡ People on MAT can’t function normally.

t

‡ People on MAT can’t take care of their children.

t

‡

‡ Historically, there have been many counselors in the field who are in recovery.

- ƒ Their beliefs about MAT influence their treatment recommendations.
- ƒ Many counselors would not recommend MAT as it conflicted with their beliefs.
- ƒ As the opioid crisis has cost so many lives, and through education and research, MAT has become a recognized pathway for recovery for many.

‡ The best way to address the mistaken beliefs is education (Westreich, 2019).

‡ Visit an opioid treatment program (OTP); talk with patients.

ƒ Access resources about evidence-based practices for

- Methadone for opioid use disorder (OUD) is highly regulated.
  - Methadone clinics are not available everywhere.
  - There are caps on the numbers of patients.
- Buprenorphine prescribing requires special certification.
  - Physicians might be deterred from offering.
  - There are caps on the number of patients.

(SAMHSA, 2012)

Without insurance, the cost of medications can be great.

‡ Naltrexone (Vivitrol) -

‡ Buprenorphine

‡ Buprenorphine is commonly diverted.

‡ Some patients with low tolerance to buprenorphine may experience effects.

‡ Most patients receiving buprenorphine treatment remain on the medication to avoid withdrawal symptoms – exactly what it is prescribed for.

‡ Preventing diversion is difficult.

(Vocci et al, 2015; Lofwall and Walsh, 2014; Bentzley et al, 2015)

t Z š [• Z o o Ç / u ‰ } Œ š v š

‡ MAT saves lives.

‡ Studies have compared MAT vs. non-MAT treatment for opioid use disorder (OUD).

f Non-MAT methods have been shown to have fatality rates as high as 20%.

f MAT methods have 0% fatalities.

(Larochelle et al, 2018; Kakko et al, 2003)

District Court Judge
6th Judicial District of Minnesota

February 24, 2020

‡ There is a need for data-driven best practices, and a move toward collecting data and evaluation.

ƒ MAT is a pressing, urgent, life-saving example.

SAMHSA
Opioid Overdose Prevention
TOOLKIT
Five Essential Steps
Information for Prescribers
Safety Advice for Patients & Family Members
Recovering From Opioid Overdose

- Change needs a champion and a team.
- Naloxone (Narcan) can be good start.
- Do we all need to be personally affected before we are willing to change our positions?
- Humanize, speak, share people’s stories.

‡ Addiction Neuroscience 101

‡

“Never doubt that a small group of thoughtful, committed citizens can change the world; indeed, it’s the only thing that ever has.”

- Margaret Mead

SAMHSA's mission is to reduce the impact of substance abuse and mental illness on America's communities.

www.samhsa.gov

1-877-SAMHSA-7 (1-877-726-4727) ● 1-800-487-4889 (TDD)

The GAINS Center focuses on expanding access to services for people with mental and/or substance use disorders who come into contact with the justice system.

https://www.samhsa.gov/gains-center

1-800-311-4246